


"Express Mail" mailing label number ET915211853US¹

Date of Deposit February 7, 2002

I hereby certify that this paper or fee
is being deposited with the United States
Postal Service "Express Mail Post Office
to Addressee" service under 37 CFR § 1.10
on the date indicated above and is
addressed to the Assistant Commissioner
for Patents, Washington, D.C. 20231

Carolyn Roberts
(Typed or printed name of person
mailing paper or fee)

(Signature of person mailing paper or fee)

PATENT
Case No. P-162-1

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE
APPLICATION FOR UNITED STATES LETTERS PATENT

INVENTORS:

Rifat Pamukcu
Gary Piazza

TITLE:

METHOD OF INHIBITING
NEOPLASTIC CELLS WITH
INDOLE DERIVATIVES

ATTORNEY:

Robert W. Stevenson
Cell Pathways, Inc.
702 Electronic Drive
Horsham, Pennsylvania 19044
(215) 706-3800

METHOD OF INHIBITING NEOPLASTIC CELLS WITH INDOLE DERIVATIVES

TECHNICAL FIELD

This invention relates to a method for the selective inhibition of neoplastic cells, for example, for the treatment or prevention of precancerous lesions or other neoplasias in mammals. This application is a continuation-in-part of U.S. patent application no. 09/007,098, which is incorporated herein by reference.

BACKGROUND OF THE INVENTION

Each year in the United States alone, untold numbers of people develop precancerous lesions, which is a form of neoplasia, as discussed below. Such lesions exhibit a strong tendency to develop into malignant tumors, or cancer. Such lesions include lesions of the breast (that can develop into breast cancer), lesions of the skin (that can develop into malignant melanoma or basal cell carcinoma), colonic adenomatous polyps (that can develop into colon cancer), and other such neoplasms. Compounds that prevent or induce the remission of existing precancerous or cancerous lesions or carcinomas would greatly reduce illness and death from cancer.

For example, approximately 60,000 people die from colon cancer, and over 150,000 new cases of colon cancer are diagnosed each year. For the American population as a whole, individuals have a six percent lifetime risk of developing colon cancer, making it the second most prevalent form of cancer in the country. Colon cancer is also prevalent in Western Europe. It is believed that increased dietary fat consumption is increasing the risk of colon cancer in Japan.

In addition, the incidence of colon cancer reportedly increases with age, particularly after the age of 40. Since the mean ages of populations in America and Western Europe are increasing, the prevalence of colorectal cancer should increase in the future.

To date, little progress has been made in the prevention and treatment of colorectal cancer, as reflected by the lack of change in the five-year survival rate over the last few decades. The only cure for this cancer is surgery at an extremely early stage. Unfortunately, most of these cancers are discovered too late for surgical cure. In many cases, the patient does not experience symptoms until the cancer has progressed to a malignant stage.

In view of these grim statistics, efforts in recent years have concentrated on colon cancer prevention. Colon cancer usually arises from pre-existing benign neoplastic growths known as polyps. Prevention efforts have emphasized the identification and removal of colonic polyps. Polyps are identified by x-ray and/or colonoscopy, and usually removed by devices associated with the colonoscope. The increased use of colon x-rays and colonoscopies in recent years has detected clinically significant precancerous polyps in four to six times the number of individuals per year that acquire colon cancer. During the past five years alone, an estimated 3.5 to 5.5 million people in the United States have been diagnosed with adenomatous colonic polyps, and it is estimated that many more people have or are susceptible to developing this condition, but are as yet undiagnosed. In fact, there are estimates that 10-12 percent of people over the age of 40 will form clinically significant adenomatous polyps.

Removal of polyps has been accomplished either with surgery or fiber-optic endoscopic polypectomy -- procedures that are uncomfortable, costly (the cost of a single polypectomy ranges between \$1,000 and \$1,500 for endoscopic treatment and more for surgery), and involve a small but significant risk of colon perforation. Overall, about \$2.5 billion is spent annually in the United States in colon cancer treatment and prevention.

In the breast, breast cancer is often treated surgically, often by radical mastectomy with its painful aftermath. Such surgery is costly, too.

As indicated above, each lesion carries with it a chance that it will develop into a cancer. The likelihood of cancer is diminished if a precancerous lesion is removed. However, many of these patients demonstrate a propensity for developing additional lesions in the future. They must, therefore, be monitored periodically for the rest of their lives for reoccurrence.

In most cases (i.e. the cases of sporadic lesion formation, e.g. so-called common sporadic polyps), lesion removal will be effective to reduce the risk of cancer. In a small percentage of cases (i.e. cases where numerous lesions form, e.g. the so-called polyposis syndromes), removal of all or part of the effected area (e.g. the colon) is indicated. For example, the difference between common sporadic polyps and polyposis syndromes is dramatic. Common sporadic polyp cases are characterized by relatively few polyps which can usually be removed leaving the colon intact. By contrast, polyposis syndrome cases can be characterized by many (e.g. hundreds or more) of polyps -- literally covering the colon in

some cases -- making safe removal of the polyps impossible short of surgical removal of the colon.

Because each lesion carries with it a palpable risk of cancerous development, patients who form many lesions (e.g. polyposis syndrome patients) invariably develop cancer if left untreated. Surgical removal of the colon is the conventional treatment in polyposis patients. Many polyposis patients have undergone a severe change in lifestyle as a result of the disfiguring surgery. Patients have strict dietary restrictions, and many must wear ostomy appliances to collect their intestinal wastes.

The search for drugs useful for treating and preventing cancer is intensive. Indeed, much of the focus of cancer research today is on the prevention of cancer because chemotherapy for cancer itself is often not effective and has severe side effects. Cancer chemoprevention is important for recovered cancer patients who retain a risk of cancer reoccurrence. Also, cancer prevention is important for people who have not yet had cancer, but have hereditary factors that place them at risk of developing cancer. With the development of new genetic screening technologies, it is easier to identify those patients with high-risk genetic factors, such as the potential for polyposis syndrome, who would greatly benefit from chemopreventative drugs. Therefore, finding such anti-cancer drugs that can be used for prolonged preventive use is of vital interest.

Known chemopreventative and chemotherapeutic drugs are believed to kill cancer cells by inducing apoptosis, sometimes referred to as "programmed cell death." Apoptosis naturally occurs in virtually all tissues of the body, and especially in self-renewing tissues such as bone marrow, immune cells, gut, liver and skin. Apoptosis plays a critical role in tissue homeostasis, that is, it ensures that the number of new cells produced are correspondingly offset by an equal number of cells that die. For example, the cells in the intestinal lining divide so rapidly that the body must eliminate cells after only three days in order to prevent the overgrowth of the intestinal lining.

Recently, scientists have realized that abnormalities of apoptosis can lead to the formation of precancerous lesions and carcinomas. Also, recent research indicates that defects in apoptosis play a major role in other diseases in addition to cancer. Consequently, compounds that modulate apoptosis could be used to prevent or control cancer, as well as used in the treatment of other diseases.

Unfortunately, even though known chemotherapeutic drugs may exhibit such

desirable apoptosis effects, most chemotherapeutic drugs have serious side effects that prohibit their long-term use, or use in otherwise healthy individuals with precancerous lesions. These side effects, which are a result of the high levels of cytotoxicity of the drugs, include hair loss, weight loss, vomiting, immune suppression and other toxicities. Therefore, there is a need to identify new drug candidates for therapy that do not have such serious side effects in humans.


In recent years, several non-steroidal anti-inflammatory drugs ("NSAIDs"), originally developed to treat arthritis, have shown effectiveness in inhibiting and eliminating colonic polyps. Polyps virtually disappear when the patients take the drug, particularly when the NSAID sulindac is administered. However, the prophylactic use of currently available NSAIDs, even in polyposis syndrome patients, is marked by severe side reactions that include gastrointestinal irritations, perforations, ulcerations and kidney toxicity. Once NSAID treatment is terminated due to such complications, the polyps return, particularly in polyposis syndrome patients.

Sulindac has been particularly well received among the NSAIDs for polyp treatment. Sulindac is a sulfoxide compound that itself is believed to be inactive as an anti-arthritic agent. The sulfoxide is reportedly converted by liver enzymes to the corresponding sulfide, which is acknowledged to be the active moiety as a prostaglandin synthesis inhibitor. The sulfide, however, is associated with the side effects of conventional NSAIDs. The sulfoxide is also known to be metabolized to a sulfone compound that has been found to be inactive as an inhibitor of prostaglandin synthesis but active as an inhibitor of precancerous lesions.

SUMMARY OF THE INVENTION

This invention includes a method of inhibiting neoplastic cells by exposing those cells to a pharmacologically effective amount of those compounds described below. Such compounds are effective in modulating apoptosis and eliminating and inhibiting the growth of neoplasias such as precancerous lesions, but are not characterized by the severe side reactions of conventional NSAIDs or other chemotherapeutics.

The compounds of that are useful in the methods of this invention include those of formula I



wherein R₁ to R₃ are independently selected from the group consisting of;

- (1) a hydrogen atom,
- (2) a lower alkyl group, a lower alkylthio group, or a lower alkoxy-lower

alkyl group, or

- (3) a lower alkyl group, an oxy group, an oxy-lower alkyl group, a lower alkyloxy group, a carbonyl group, a lower alkenyl group, an optionally-substituted imino group, a lower alkylimino group optionally substituted at its nitrogen atom, a thio-lower alkyl group, or a lower alkylthio group; or to each group in (3), bonded is an aryl group or a heterocyclic group, or each group in (3) is substituted by an aryl group or a heterocyclic group; and said aryl or heterocyclic group may be further substituted by a halogen atom, a nitro group, a lower alkylamino group, an acylamino group, a lower alkyl group, a lower alkoxy group, a halo-lower alkyl group, a lower cycloalkyl group, or an aryl, heterocyclic, aryl-lower alkyl, heterocyclic-lower alkyl, aryl-lower alkyloxy, heterocyclic-lower alkyloxy, aryl-lower alkenyl or heterocyclic-lower alkenyl group optionally substituted by any of a halogen atom or a lower alkyl group, with the proviso that R₁ to R₃ must not be simultaneously all hydrogen atoms;

R₄ is selected from the group consisting of a hydrogen atom, or a lower alkyl group;

R₅ is selected from the group consisting of a carboxyl group, an esterified

carboxyl group, or an amidated carboxyl group.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

As indicated above, this invention relates to a method for inhibiting neoplasia, particularly cancerous and precancerous lesions by exposing the affected cells to a compound of Formula I above.

Preferably, such compounds are administered without therapeutic amounts of an NSAID.

The present invention is also a method of treating mammals with precancerous lesions by administering a pharmacologically effective amount of an enterically coated pharmaceutical composition that includes compounds of this invention.

Also, the present invention is a method of inhibiting the growth of neoplastic cells by exposing the cells to an effective amount of compounds of Formula I, wherein R_1 through R_3 are defined as above.

In still another form, the invention is a method of inducing apoptosis in human cells by exposing those cells to an effective amount of compounds of Formula I to those cells sensitive to such a compound.

As used herein, the term "precancerous lesion" includes syndromes represented by abnormal neoplastic, including dysplastic, changes of tissue.

Examples include adenomatous growths in colonic, breast or lung tissues, or conditions such as dysplastic nevus syndrome, a precursor to malignant melanoma of the skin. Examples also include, in addition to dysplastic nevus syndromes, polyposis syndromes, colonic polyps, precancerous lesions of the cervix (i.e., cervical dysplasia), prostatic dysplasia, bronchial dysplasia, breast, bladder and/or skin and related conditions (e.g., actinic keratosis), whether the lesions are clinically identifiable or not.

As used herein, the term "carcinomas" refers to lesions that are cancerous. Examples include malignant melanomas, breast cancer, and colon cancer.

As used herein, the term "neoplasm" refers to both precancerous and cancerous lesions.

Compounds useful in the methods of this invention may be formulated into compositions together with pharmaceutically acceptable carriers for oral administration in

solid or liquid form, or for rectal administration, although carriers for oral administration are most preferred.

Pharmaceutically acceptable carriers for oral administration include capsules, tablets, pills, powders, troches and granules. In such solid dosage forms, the carrier can comprise at least one inert diluent such as sucrose, lactose or starch. Such carriers can also comprise, as is normal practice, additional substances other than diluents, e.g., lubricating agents such as magnesium stearate. In the case of capsules, tablets, troches and pills, the carriers may also comprise buffering agents. Carriers such as tablets, pills and granules can be prepared with enteric coatings on the surfaces of the tablets, pills or granules.

Alternatively, the enterically coated compound can be pressed into a tablet, pill, or granule, and the tablet, pill or granules for administration to the patient. Preferred enteric coatings include those that dissolve or disintegrate at colonic pH such as shellac or Eudraget S.

Pharmaceutically acceptable carriers include liquid dosage forms for oral administration, e.g. pharmaceutically acceptable emulsions, solutions, suspensions, syrups and elixirs containing inert diluents commonly used in the art, such as water. Besides such inert diluents, compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, and sweetening, flavoring and perfuming agents.

Pharmaceutically acceptable carriers for rectal administration are preferably suppositories that may contain, in addition to the compounds of Formula I, excipients such as cocoa butter or a suppository wax.


The pharmaceutically acceptable carrier and compounds of this invention are formulated into unit dosage forms for administration to a patient. The dosage levels of active ingredient (i.e. compounds of this invention) in the unit dosage may be varied so as to obtain an amount of active ingredient effective to achieve lesion-eliminating activity in accordance with the desired method of administration (i.e., oral or rectal). The selected dosage level therefore depends upon the nature of the active compound administered, the route of administration, the desired duration of treatment, and other factors. If desired, the unit dosage may be such that the daily requirement for active compound is in one dose, or divided among multiple doses for administration, e.g., two to four times per day.

The pharmaceutical compositions of this invention are preferably packaged in a container (e.g. a box or bottle, or both) with suitable printed material (e.g. a package insert) containing indications, directions for use, etc.

The amount of the derivative to be used varies, depending on the age and the condition of the patients, on the type and the condition of the diseases, and on the type of the derivative. In general, however, the derivative may be administered in an amount between 1 and 100 mg/kg for oral administration, or between 0.1 and 10 mg/kg for intramuscular or intravenous administration, once to four times a day. In practice, the physician will determine the actual dosing regimen that will be most suitable for an individual patient and it will vary with the age, weight and response of the particular patient. The above dosages are believed to be exemplary of the average case, but there may be individual instances in which higher or lower dosage ranges may be merited, and such are within the scope of this invention.


Methods for producing the indole derivatives useful in this invention are reported in WO 98/145530 (which is incorporated herein by reference), and are set forth below.

As in the reaction formula (a), a compound (1) is processed with a strong base such as sodium hydride, lithium diisopropylamide or the like, and then reacted with a chloride, bromide, iodide, toluenesulfonate or methane sulfonate of $R^{1'a}$ to produce a compound (2). Alternatively, the compound (2) may also be produced, using an inorganic or organic base such as typically potassium carbonate, sodium carbonate, lithium carbonate, sodium hydrogencarbonate, potassium hydrogencarbonate, pyridine or triethylamine. In this case, where $R^{1'a}$ in (1) is a hydrogen atom, the compound (1) reacts with the reactant also at its 3-position.




wherein R^{1a} , $R^{1'a}$ and R^{2a} each represent 1) a hydrogen atom, 2) a lower alkyl group, a lower alkylthio group, or a lower alkoxy-lower alkyl group, or 3) a lower alkyl group, an oxy group, an oxy-lower alkyl group, a lower alkyloxy group, a carbonyl group, a lower alkenyl group, an optionally-substituted imino group, a lower alkylimino group optionally substituted at its nitrogen atom, a thio-lower alkyl group, or a lower alkylthio group; to each group in 3), bonded is an aryl group or a heterocyclic group, or each group in 3) is substituted by an aryl group or a heterocyclic group, and said aryl or heterocyclic group may be further substituted by any of a halogen atom, a nitro group, a lower alkylamino group, an acylamino group, a lower alkyl group, a lower alkoxy group, a halo-lower alkyl group, a lower cycloalkyl group, or an aryl, heterocyclic, aryl-lower alkyl, heterocyclic-lower alkyl, aryl-lower alkyloxy, heterocyclic-lower alkyloxy, aryl-lower alkenyl or heterocyclic-lower alkenyl group optionally substituted by any of a halogen atom or a lower alkyl group; but R^{1a} , $R^{1'a}$ and R^{2a} must not be all hydrogen atoms at the same time; R^{3a} represents an alkyl group; and R^{4a} represents a hydrogen atom or a lower alkyl group.

The compound (2) is hydrolyzed with a base such as lithium hydroxide, sodium hydroxide, potassium hydroxide or the like to produce a compound (3) (reaction formula (b)).




wherein R^{1a} , $R^{1'a}$, R^{2a} , R^{3a} and R^{4a} have the same meanings as above.

The compound (3) is processed with a carboxyl activator such as carbonyldiimidazole, 1-(3-(dimethylamino)propyl-3-ethylcarbodiimide, dicyclohexylcarbodiimide or the like, and then reacted with a sulfonamide in the presence of a base to produce a compound (4) (reaction formula (c)). In this case, where the sulfonamide used has a reactive substituent, sultams may be produced through cyclization.




wherein R^{1a} , $\text{R}^{1'a}$, R^{2a} and R^{4a} have the same meanings as above; R^{6a} represents an alkyl, alkenyl, lower alkoxy-lower alkyl, aryl, heterocyclic or lower cycloalkyl group, which may be optionally substituted by at least one selected from a halogen atom, a lower alkoxy group, a lower alkyl group, a lower alkenyl group, a lower cycloalkyl group, a nitro group, an aryl group, a heterocyclic group, an arylazo group, a halo-lower alkyl group, a lower alkylaryl group and a lower alkoxyaryl group; and the nitrogen atom of the sulfonamido group in the formula may release the hydrogen atom bonding thereto to form a ring along with R^{6a}

As in the reaction formula (a), a compound (6) can be produced from the compound (5) (reaction formula (d)).




wherein $R^{1'a}$, R^{2a} and R^{4a} have the same meanings as above; R^{1b} represents a lower alkanoyl group, an arylcarbonyl group, or a heterocyclic-carbonyl group, and said arylcarbonyl or heterocyclic-carbonyl group may be further substituted by any of a halogen atom, a nitro group, a lower alkylamino group, an acylamino group, a lower alkyl group, a lower alkoxy group, a halo-lower alkyl group, a lower cycloalkyl group, or an aryl, heterocyclic, aryl-lower alkyl, heterocyclic-lower alkyl, aryl-lower alkyloxy, heterocyclic-lower alkyloxy, aryl-lower alkenyl or heterocyclic-lower alkenyl group optionally substituted by any of a halogen atom or a lower alkyl group; and R^{3a} represents a lower alkyl group.

In the reaction formula (b), a compound (7) can be produced from the compound (6) (reaction formula (e)).




wherein $R^{1'a}$, R^{2a} , R^{1b} and R^{4a} have the same meanings as above; and R^{3a} represents a lower alkyl group.

In the reaction formula (c), a compound (S) can be produced from the compound (7) (reaction formula (f)). In this case, where the sulfonamide used has a reactive substituent, sultams may be produced through cyclization.




wherein $R^{1'a}$, R^{2a} , R^{1b} , R^{4a} and R^{6a} have the same meanings as above.

In the reaction formula (g), a compound (9) is reacted with a chloride, bromide or iodide of $R^{1'a}$ in the presence of silver(I) oxide, whereby the compound (9) is converted into a compound (10). Presence of sodium iodide or potassium iodide in the reaction will improve the result. Alternatively, the compound (9) may be reacted in the same manner as above but in the presence of potassium hydrogentartrate or sodium hydrogentartrate to obtain the compound (10). Presence of sodium iodide or potassium iodide in the reaction will also improve the result. In place of the chloride, bromide or iodide used herein, a corresponding toluenesulfonate or methanesulfonate may also be used to obtain the compound (10).




wherein R^{1a} , $R^{1'a}$, R^{2a} and R^{4a} have the same meanings as above; and R^{3a} represents an alkyl group.

In the reaction formula (b), a compound (11) is produced from the compound (10) (reaction formula (h)).



wherein R^{1a} , $R^{1'a}$, R^{2a} and R^{4a} have the same meanings as above; and R^{3a} represents an alkyl group.


In the reaction formula (c), a compound (12) can be produced from the compound (11) (reaction formula (i)). In this case, where the sulfonamide used has a reactive substituent, sultams may be produced through cyclization.



wherein R^{1a} , $R^{1'a}$, R^{2a} , R^{4a} and R^{6a} have the same meanings as above.

In the reaction formula (j), the compound (9) is reacted with a lower alkanoyl


chloride or bromide, or arylcarbonyl chloride or bromide, or a heterocyclic-carbonyl chloride or bromide (said arylcarbonyl chloride or bromide, or heterocyclic-carbonyl chloride or bromide may be substituted by any of a halogen atom, a nitro group, a lower alkylamino group, an acylamino group, a lower alkyl group, a lower alkoxy group, a halo-lower alkyl group, a lower cycloalkyl group, or an aryl, heterocyclic, aryl-lower alkyl, heterocyclic-lower alkyl, aryl-lower alkyloxy, heterocyclic-lower alkyloxy, aryl-lower alkenyl or heterocyclic-lower alkenyl group optionally substituted by any of a halogen atom or a lower alkyl group), in the presence of a strong Lewis acid such as aluminium chloride, tin(IV) chloride, tin(II) chloride, iron(III) chloride, boron trifluoride, zirconia sulfate or the like, whereby the compound (9) is converted into a compound (13).



wherein R^{1a} , R^{2a} , R^{1b} and R^{4a} have the same meanings as above; and R^{3a} represents an alkyl group.

In the reaction formula (k), the compound (13) is converted into a compound (14), by processing it with a metal hydride such as typically sodium borohydride or borane-tetrahydrofuran complex. The compound (14) is converted into a compound (15), by processing it with sodium borohydride in the presence of trifluoroacetic acid, trifluoromethanesulfonic acid or sulfuric acid. Alternatively, the compound (14) may


also be converted into the compound (15), by processing it with a trialkylsilane such as typically triethylsilane in the presence of trifluoroacetic acid. If the compound (13) is processed with a trialkylsilane such as typically triethylsilane in the presence of trifluoroacetic acid, it may be directly converted into the compound (15) in one-step reaction.



wherein $\text{R}^{1\text{a}}$, $\text{R}^{1'\text{a}}$, $\text{R}^{2\text{a}}$ and $\text{R}^{4\text{a}}$ have the same meanings as above; $\text{R}^{3\text{a}}$ represents an alkyl group; $\text{R}^{1\text{c}}$ represents a lower alkylhydroxymethyl group, an arylhydroxymethyl group, or a heterocyclic-hydroxymethyl group, and said arylhydroxymethyl group or heterocyclic-hydroxymethyl group may be substituted by any of a halogen atom, a nitro group, a lower alkylamino group, an acylamino group, a lower alkyl group, a lower alkoxy group, a halo-lower alkyl group, a lower cycloalkyl group, or an aryl, heterocyclic, aryl-lower alkyl, heterocyclic-lower alkyl, aryl-lower alkyloxy, heterocyclic-lower alkyloxy, aryl-lower alkenyl or heterocyclic-lower alkenyl group


optionally substituted by any of a halogen atom or a lower alkyl group.

In the reaction formula (k), a compound (18) can be produced from a compound (16) via a compound (17) (reaction formula (l)).



wherein R^{1a} , $R^{1'a}$, R^{1b} , R^{1c} , R^{2a} and R^{4a} have the same meanings as above.

In the reaction formula (k), a compound (21) can be produced from a compound (19) via a compound (20) (reaction formula (m)).




wherein R^{1a} , $R^{1'a}$, R^{1b} , R^{1c} , R^{2a} , R^{4a} and R^{6a} have the same meanings as above.

The compound (13) may be reacted with an aldehyde in the presence of

trifluoroacetic acid and a trialkylsilane such as typically triethylsilane, whereby it is converted into the compound (18) via the compound (17). Alternatively, the compound (13) may be reacted with hydrazine and an alkali such as typically sodium hydroxide or potassium hydroxide, whereby it is converted directly into the compound (18).

The compound (1) reportedly can be produced according to the methods described in "Journal of Medicinal Chemistry, 1992, 35, 2419". "US-5212195". "US-4894386", "EP-0242167". Specifically, a compound (22) is reacted with an N,N-dialkylformamide such as typically N,N-dimethylformamido-dimethyl acetal to produce a compound (23). Where R^{2a} in (23) is 1) a lower alkyl group, a lower alkylthio group, or a lower alkoxy-lower alkyl group, or 2) a lower alkyl group, an oxy group, an oxy-lower alkyl group, a lower alkyloxy group, a carbonyl group, a lower alkenyl group, an optionally-substituted imino group, a lower alkylimino group optionally substituted at its nitrogen atom, a thio-lower alkyl group, or a lower alkylthio group (to each group in 2), bonded is an aryl group or a heterocyclic group, or each group in 2) is substituted by an aryl group or a heterocyclic group, and said aryl or heterocyclic group may be further substituted by any of a halogen atom, a nitro group, a lower alkylamino group, an acylamino group, a lower alkyl group, a lower alkoxy group, a halo-lower alkyl group, a lower cycloalkyl group, or an aryl, heterocyclic, aryl-lower alkyl, heterocyclic-lower alkyl, aryl-lower alkyloxy, heterocyclic-lower alkyloxy, aryl-lower alkenyl or heterocyclic-lower alkenyl group optionally substituted by any of a halogen atom or a lower alkyl group), the compound (23) may be reacted with a lower alkanoyl chloride or bromide to give a compound (24), which is then reduced at its nitro group in the presence of a hydrogenation catalyst such as typically palladium-carbon, in a hydrogen atmosphere, to produce the compound (1) (reaction formula (n)). Apart from this mode, the nitro group may be reduced by any other method of, for example, 1) reduction with

reduced iron and zinc, 2) reduction with sodium hydrosulfite, 3) reduction with formic acid or ammonium formate in the presence of a transition metal catalyst such as typically palladium-carbon, or 4) reduction with nickel.




(n)

wherein R^{1a} and R^{4a} have the same meanings as above; R^{2a} represents 1) a lower alkyl group, a lower alkylthio group, or a lower alkoxy-lower alkyl group, or 2) a lower alkyl group, an oxy group, an oxy-lower alkyl group, a lower alkyloxy group, a carbonyl group, a lower alkenyl group, an optionally-substituted imino group, a lower alkylimino group optionally substituted at its nitrogen atom, a thio-lower alkyl group, or a lower alkylthio group, to each group in 2), bonded is an aryl group or a heterocyclic group, or each group in 2) is substituted by an aryl group or a heterocyclic group, and said aryl or heterocyclic group may be further substituted by any of a halogen atom, a nitro group, a lower alkylamino group, an acylamino group, a lower alkyl group, a lower alkoxy

group, a halo-lower alkyl group, a lower cycloalkyl group, or an aryl, heterocyclic, aryl-lower alkyl, heterocyclic-lower alkyl, aryl-lower alkyloxy, heterocyclic-lower alkyloxy, aryl-lower alkenyl or heterocyclic-lower alkenyl group optionally substituted by any of a halogen atom or a lower alkyl group; R^{2a} represents an alkyl group; and R' represents a lower alkyl group.


A compound (25) which corresponds to the compound (1) where R^{3a} is a hydrogen atom can be produced by reducing a compound (23) at its nitro group in the presence of a hydrogenation catalyst such as typically palladium-carbon, in a hydrogen atmosphere (reaction formula (o)). Apart from this mode, the nitro group may be reduced by any other method of, for example, 1) reduction with reduced iron and zinc, 2) reduction with sodium hydrosulfite, 3) reduction with formic acid or ammonium formate in the presence of a transition metal catalyst such as typically palladium-carbon, or 4) reduction with nickel.



wherein R^{1a} , R^{4a} and R' have the same meanings as above; and R^{3a} represents an alkyl group.

In the reaction formula (p), a compound (26), which corresponds to the compound (1) where R^{1a} is a lower alkyl group and R^{2a} is a hydrogen atom, can be converted into a compound (29) which corresponds to the compound (1) where R^{1a} is a

lower alkyl group and R^{2a} is a lower alkyl group or an aryl-lower alkyl group (said aryl-lower alkyl group may be substituted by any of a halogen atom, a nitro group, a lower alkylamino group, an acylamino group, a lower alkyl group, a lower alkoxy group, a halo-lower alkyl group, a lower cycloalkyl group, or an aryl, heterocyclic, aryl-lower alkyl, heterocyclic-lower alkyl, aryl-lower alkyloxy, heterocyclic-lower alkyloxy, aryl-lower alkenyl or heterocyclic-lower alkenyl group optionally substituted by any of a halogen atom or a lower alkyl group). Specifically, the compound (26) can be converted into a compound (27), like in the reaction formula (j). The compound (27) can be converted into the compound (29) via a compound (28), like in the reaction formula (k).



(p)

wherein R^{1a} represents a lower alkyl group; R^{2b} represents a lower alkyl group or an

aryl-lower alkyl group (said aryl-lower alkyl group may be substituted by any of a halogen atom, a nitro group, a lower alkylamino group, an acylamino group, a lower alkyl group, a lower alkoxy group, a halo-lower alkyl group, a lower cycloalkyl group, or an aryl, heterocyclic, aryl-lower alkyl, heterocyclic-lower alkyl, aryl-lower alkyloxy, heterocyclic-lower alkyloxy, aryl-lower alkenyl or heterocyclic-lower alkenyl group optionally substituted by any of a halogen atom or a lower alkyl group); R^{3a} represents a lower alkyl group; R^{2c} represents a lower alkyl group or an aryl-lower alkyl group (said aryl-lower alkyl group may be substituted by any of a halogen atom, a nitro group, a lower alkylamino group, an acylamino group, a lower alkyl group, a lower alkoxy group, a halo-lower alkyl group, a lower cycloalkyl group, or an aryl, heterocyclic, aryl-lower alkyl, heterocyclic-lower alkyl, aryl-lower alkyloxy, heterocyclic-lower alkyloxy, aryl-lower alkenyl or heterocyclic-lower alkenyl group optionally substituted by any of a halogen atom or a lower alkyl group); and R^{4a} has the same meaning as above.

The intermediates in each reaction noted above may optionally be purified in any ordinary purification for chemical synthesis, for example, through recrystallization, column chromatography, thin-layer chromatography, high-performance liquid chromatography or the like, prior to the next step where they are processed. The final products of the compounds of the invention may also be purified, if desired, in any ordinary purification for organic chemistry, for example, through recrystallization, column chromatography, thin-layer chromatography, high-performance liquid chromatography or the like. Identification of the compounds may be effected through NMR spectrometric analysis, mass spectrometric analysis, IR spectrometric analysis, elementary analysis, melting point measurement, etc.

Unless otherwise specifically indicated, "lower" is meant to indicate groups having at most 8 carbon atoms.

The alkyl group is meant to have from 1 to 20 carbon atoms. This may be a linear or branched alkyl group, including, for example, a methyl group, an ethyl group, an n-propyl group, an i-propyl group, an n-butyl group, an i-butyl group, a sec-butyl group, a t-butyl group, an n-pentyl group, an i-pentyl group, a sec-pentyl group, a 2,2-dimethylpentyl group, a 2-methylbutyl group, an n-hexyl group, a 1-methylpentyl group, a 2-methylpentyl group, a 3-methylpentyl group, a 4-methylpentyl group, a 1-ethylbutyl group, a 2-ethylbutyl group, a 1,1-dimethylbutyl group, a 2,2-dimethylbutyl group, a 3,3-dimethylbutyl group, a 1-ethyl-1-methylpropyl group, an n-heptyl group, a 1-methylhexyl group, a 2-methylhexyl group, a 3-methylhexyl group, a 4-methylhexyl group, a 5-methylhexyl group, a 1-ethylpentyl group, a 2-ethylpentyl group, a 1,1-dimethylpentyl group, a 2,2-dimethylpentyl group, a 3,3-dimethylpentyl group, an n-octyl group, a 1-methylheptyl group, a 2-methylheptyl group, a 3-methylheptyl group, a 4-methylheptyl group, a 5-methylheptyl group, a 6-methylheptyl group, a 1-ethylhexyl group, a 2-ethylhexyl group, a 1,1-dimethylhexyl group, a 2,2-dimethylhexyl group, a 3,3-dimethylhexyl group, an n-nonyl group, a 1-methyloctyl group, a 2-methyloctyl group, a 3-methyloctyl group, a 4-methyloctyl group, a 5-methyloctyl group, a 6-methyloctyl group, a 7-methyloctyl group, a 1-ethylheptyl group, a 2-ethylheptyl group, a 1,1-dimethylheptyl group, a 2,2-dimethylheptyl group, a 3,3-dimethylheptyl group, an n-decyl group, a 1-methylnonyl group, a 2-methylnonyl group, a 3-methylnonyl group, a 4-methylnonyl group, a 1-ethyloctyl group, a 2-ethyloctyl group, an n-undecyl group, an n-dodecyl group, an n-tridecyl group, an n-tetradecyl group, an n-pentadecyl group, an n-hexadecyl group, an n-octadecyl group, etc. Preferred are alkyl groups each having from 3 to 8 carbon atoms.

The alkenyl group is meant to have from 2 to 20 carbon atoms. This may be a linear or branched alkenyl group, including, for example, a vinyl group, a 1-propenyl

group, a 2-propenyl group, a 1-butenyl group, a 2-butenyl group, a 3-butenyl group, a 1-methyl-1-propenyl group, a 2-methyl-1-propenyl group, a 1-methyl-2-propenyl group, a 2-methyl-2-propenyl group, a 1-pentenyl group, a 2-pentenyl group, a 3-pentenyl group, a 4-pentenyl group, a 1-methyl-1-butenyl group, a 2-methyl-1-butenyl group, a 3-methyl-1-butenyl group, a 2-methyl-2-butenyl group, a 3-methyl-2-butenyl group, a 2-methyl-3-butenyl group, a 3-methyl-3-butenyl group, a 1-hexenyl group, a 2-hexenyl group, a 3-hexenyl group, a 4-hexenyl group, a 5-hexenyl group, a 2-methyl-1-pentenyl group, a 3-methyl-1-pentenyl group, a 4-methyl-1-pentenyl group, a 1-heptenyl group, a 1-octenyl group, a 1-nonenyl group, a 1-decenyl group, a 1-undecenyl group, a 1-dodecenyl group, a 1-tridecenyl group, a 1-tetradecenyl group, a 1-pentadecenyl group, a 1-hexadecenyl group, a 1-octadecenyl group, etc. Preferred are alkenyl groups each having from 3 to 8 carbon atoms.

As preferred examples of the alkenyl group, mentioned are linear or branched alkenyl groups including a vinyl group, a 1-propenyl group, a 2-propenyl group, a 1-butenyl group, a 2-butenyl group, a 3-butenyl group, a 1-methyl-1-propenyl group, a 2-methyl-1-propenyl group, a 1-methyl-2-propenyl group, a 2-methyl-2-propenyl group, a 1-pentenyl group, a 2-pentenyl group, a 3-pentenyl group, a 4-pentenyl group, a 1-methyl-1-butenyl group, a 2-methyl-1-butenyl group, a 3-methyl-1-butenyl group, a 2-methyl-2-butenyl group, a 3-methyl-2-butenyl group, a 2-methyl-3-butenyl group, a 3-methyl-3-butenyl group, a 1-hexenyl group, a 2-hexenyl group, a 3-hexenyl group, a 4-hexenyl group, a 5-hexenyl group, a 2-methyl-1-pentenyl group, a 3-methyl-1-pentenyl group, a 4-methyl-1-pentenyl group, etc. More preferred are those each having from 2 to 4 carbon atoms.

The halogen atom includes a fluorine atom, a chlorine atom, a bromine atom, and an iodine atom. Preferred are a fluorine atom, a chlorine atom and a bromine atom.

The halo-lower alkyl group is a linear or branched alkyl group having at most 8 carbon atoms and substituted by any of a fluorine atom, a chlorine atom, a bromine atom and an iodine atom, preferably substituted by any of a fluorine atom, a chlorine atom and a bromine atom and having at most 8 carbon atoms, more preferably having from 1 to 3 carbon atoms. For example, it includes a fluoromethyl group, a difluoromethyl group, a trifluoromethyl group, a chloromethyl group, a dichloromethyl group, a trichloromethyl group, a bromomethyl group, a dibromomethyl group, a tribromomethyl group, a 1-fluoroethyl group, a 1-chloroethyl group, a 1-bromoethyl group, a 2-fluoroethyl group, a 2-chloroethyl group, a 2-bromoethyl group, a 1,2-difluoroethyl group, a 1,2-dichloroethyl group, a 1,2-dibromoethyl group, a 2,2,2-trifluoroethyl group, a heptafluoroethyl group, a 1-fluoropropyl group, a 1-chloropropyl group, a 1-bromopropyl group, a 2-fluoropropyl group, a 2-chloropropyl group, a 2-bromopropyl group, a 3-fluoropropyl group, a 3-chloropropyl group, a 3-bromopropyl group, a 1,2-difluoropropyl group, a 1,2-dichloropropyl group, a 1,2-dibromopropyl group, a 2,3-difluoropropyl group, a 2,3-dichloropropyl group, a 2,3-dibromopropyl group, a 3,3,3-trifluoropropyl group, a 2,2,3,3,3-pentafluoropropyl group, a 2-fluorobutyl group, a 2-chlorobutyl group, a 2-bromobutyl group, a 4-fluorobutyl group, a 4-chlorobutyl group, a 4-bromobutyl group, a 4,4,4-trifluorobutyl group, a 2,2,3,3,4,4,4-heptafluorobutyl group, a perfluorobutyl group, a 2-fluoropentyl group, a 2-chloropentyl group, a 2-bromopentyl group, a 5-fluoropentyl group, a 5-chloropentyl group, a 5-bromopentyl group, a perfluoropentyl group, a 2-fluorohexyl group, a 2-chlorohexyl group, a 2-bromohexyl group, a 6-fluorohexyl group, a 6-chlorohexyl group, a 6-bromohexyl group, a perfluorohexyl group, a 2-fluoroheptyl group, a 2-chloroheptyl group, a 2-bromoheptyl group, a 7-fluoroheptyl group, a 7-chloroheptyl group, a 7-bromoheptyl group, a perfluoroheptyl group, etc.

The lower alkoxy group is a linear or branched alkyloxy group having at most 6 carbon atoms. For example, it includes a methoxy group, an ethoxy group, an n-propyloxy group, an i-propyloxy group, an n-butyloxy group, an i-butyloxy group, a sec-butyloxy group, a t-butyloxy group, an n-pentyloxy group, an i-pentyloxy group, a sec-pentyloxy group, a 2,2-dimethylpropyloxy group, a 2-methylbutoxy group, an n-hexyloxy group, an i-hexyloxy group, a t-hexyloxy group, a sec-hexyloxy group, a 2-methylpentyloxy group, a 3-methylpentyloxy group, a 1-ethylbutyloxy group, a 2-ethylbutyloxy group, a 1,1-dimethylbutyloxy group, a 2,2-dimethylbutyloxy group, a 3,3-dimethylbutyloxy group, a 1-ethyl-1-methylpropyloxy group, etc. Preferred are a methoxy group, an ethoxy group, an n-propyloxy group, an i-propyloxy group, an n-butyloxy group, an i-butyloxy group, a sec-butyloxy group, a t-butyloxy group, etc. More preferred are those each having from 1 to 3 carbon atoms.

The lower alkylthio group is meant to indicate a thio group to which is bonded a lower alkyl group.

The lower alkoxy-lower alkyl group is meant to indicate a lower alkyl group substituted by a lower alkoxy group.

The oxy-lower alkyl group is meant to indicate a lower alkyl group substituted by an oxy group.

The lower alkyloxy group is meant to indicate an oxy group to which is bonded a lower alkyl group.

The optionally-substituted imino group is meant to indicate an imino group which may be optionally substituted by a lower alkyl group or the like.

The lower alkylimino group optionally substituted at its nitrogen atom is meant to indicate an imino group to which is bonded a lower alkyl group and in which the nitrogen atom may be optionally substituted by a lower alkyl group or the like.

The thio-lower alkyl group is meant to indicate a lower alkyl group substituted by a thio group.

The lower alkylthio group is meant to indicate a thio group to which is bonded a lower alkyl group.

The lower cycloalkyl group is a cycloalkyl group having from 3 to 7 carbon atoms. Preferably, it includes a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, etc. More preferred are a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, and a cyclohexyl group.

Preferred examples of the "esterified carboxyl group" are mentioned below.

As preferred examples of the ester moiety in the esterified carboxyl group, mentioned are a lower alkyl ester (e.g., methyl ester, ethyl ester, propyl ester, isopropyl ester, butyl ester, isobutyl ester, tert-butyl ester, pentyl ester, hexyl ester, etc.). The lower alkyl ester may have at least one suitable substituent. As its examples, mentioned are a lower alkanoyloxy-(lower) alkyl ester [e.g., acetoxymethyl ester, propionyloxymethyl ester, butyryloxymethyl ester, valeryloxymethyl ester, pivaloyloxymethyl ester, hexanoyloxymethyl ester, 1- (or 2-)acetoxylethyl ester, 1- (or 2- or 3-)acetoxypentyl ester, 1-(or 2-, 3- or 4-)acetoxypentyl ester, 1-(or 2-) propionyloxyethyl ester, 1- (or 2- or 3-)propionyloxypropyl ester, 1- (or 2-) butyryloxyethyl ester, 1- (or 2-)isobutyryloxyethyl ester, 1- (or 2-)pivaloyloxyethyl ester, 1- (or 2-)hexanoyloxyethyl ester, isobutyryloxymethyl ester, 2-ethylbutyryloxymethyl ester, 3,3-dimethylbutyryloxymethyl ester, 1- (or 2-)pentanoyloxyethyl ester, etc.], a lower alkanesulfonyl-(lower) alkyl ester (e.g., 2-mesylethyl ester, etc.), a mono- (or di- or tri-)halo-(lower) alkyl ester (e.g., 2-iodoethyl ester, 2,2,2-trichloroethyl ester, etc.); a lower alkoxycarbonyloxy-(lower) alkyl ester [e.g., methoxycarbonyloxymethyl ester, ethoxycarbonyloxymethyl ester,

propoxycarbonyloxymethyl ester, tert-butoxycarbonyloxymethyl ester, 1- (or 2-)methoxycarbonyloxyethyl ester, 1- (or 2-)ethoxycarbonyloxyethyl ester, 1- (or 2-)isopropoxycarbonyloxyethyl ester, etc.], a phthalidylidene-(lower) alkyl ester, or a (5-lower alkyl-2-oxo-1,3-dioxol-4-yl)-(lower) alkyl ester [e.g., (5-methyl-2-oxo-1,3-dioxol-4-yl)methyl ester, (5-ethyl-2-oxo-1,3-dioxol-4-yl)methyl ester, (5-propyl-2-oxo-1,3-dioxol-4-yl)ethyl ester, etc.]; a lower alkenyl ester (e.g., vinyl ester, allyl ester, etc.); a lower alkynyl ester (e.g., ethynyl ester, propynyl ester, etc.); an aryl-(lower) alkyl ester optionally having at least one suitable substituent (e.g., benzyl ester, 4-methoxybenzyl ester, 4-nitrobenzyl ester, phenethyl ester, trityl ester, benzhydryl ester, bis(methoxyphenyl)methyl ester, 3,4-dimethoxybenzyl ester, 4-hydroxy-3,5-di-tert-butylbenzyl ester, etc.); an aryl ester optionally having at least one suitable substituent (e.g., phenethyl ester, 4-chlorophenyl ester, tolyl ester, tert-butylphenyl ester, xylyl ester, mesityl ester, cumenyl ester, etc.); a phthalidyl ester, etc.

As preferred examples of the protected, esterified carboxyl group which is defined as above, mentioned are a lower alkoxycarbonyl group, and a phenyl (or nitrophenyl)-(C₁₋₄) alkoxycarbonyl group. The most preferred are methoxycarbonyl, ethoxycarbonyl and benzyloxycarbonyl groups.

Preferred examples of the "amidated carboxyl group" are mentioned below.

A carbamoyl group;

A mono- or di-lower alkylcarbamoyl group (in which the alkyl group may be any of those mentioned hereinabove) [e.g., methylcarbamoyl, dimethylcarbamoyl, isopropylcarbamoyl, n-butylcarbamoyl, t-butylcarbamoyl, N-methyl-N-(pyridylmethyl)carbamoyl, etc.];

An aryl-lower alkylcarbamoyl group (in which the aryl group and the alkyl group may be any of those mentioned hereinabove) [e.g., benzylcarbamoyl, 3,4-

methylenedioxybenzylcarbamoyl, diaminobenzylcarbamoyl, phenethylcarbamoyl];

A cyclo-lower alkylcarbamoyl group having from 3 to 7 carbon atoms (in which the cyclo-lower alkyl group may be any of those mentioned hereinabove) [e.g., cyclopropylcarbamoyl, cyclobutylcarbamoyl, cyclopentylcarbamoyl, cyclohexylcarbamoyl, etc.];

An arylcarbamoyl group (in which the aryl group may be any of those mentioned hereinabove) [e.g., phenylcarbamoyl, naphthylcarbamoyl, etc.];

A heterocyclic carbamoyl group (in which the heterocyclic group may be any of those mentioned hereinabove) [e.g., thiazolylcarbamoyl, thiadiazolylcarbamoyl, pyridylcarbamoyl, triazolylcarbamoyl, tetrazolylcarbamoyl, N-methyl-N-pyridylcarbamoyl, morpholinocarbamoyl, etc.];

A heterocyclic-lower alkylcarbamoyl group (in which the heterocyclic-lower alkyl group may be any of those mentioned hereinabove) [e.g., morpholinoethylcarbamoyl, pyridylmethylcarbamoyl, methylenedioxybenzylcarbamoyl, etc.];

An N-disubstituted carbamoyl group in which the nitrogen atom is one member of atoms constituting a nitrogen-containing hetero-ring (e.g., morpholinocarbonyl, thiomorpholinocarbonyl, 1-perhydroazepinylcarbonyl, 1,1,-dioxothiazolidinecarbonyl, piperidinocarbonyl, 1-piperazinylcarbonyl, 4-(2-hydroxyethyl)-1-piperazinylcarbonyl, 4-methyl-1-piperazinylcarbonyl, carboxypyrrolidinocarbonyl, etc.);

A substituted sulfonylcarbamoyl group, etc.

The substituents for the substituted sulfonylcarbamoyl group include, for example, an alkyl group, an alkenyl group, a halo-lower alkyl group, an aryl-lower alkyl group, a hydroxy-lower alkyl group, a tri-lower alkylsilyl-lower alkyl group, a lower alkoxy-lower alkyl group, a lower alkylthio-lower alkyl group, a heterocyclic group and

an aryl group each having at most 20 carbon atoms, such as those mentioned above. The aryl group may optionally be substituted by any of a halogen atom, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group, a nitro group and the like, such as those mentioned above. As specific examples of the substituted sulfonylcarbonyl group, concretely mentioned are a naphthalenesulfonylcarbamoyl group, a benzenesulfonylcarbamoyl group, a nitrobenzenesulfonylcarbamoyl group, a trihalobenzenesulfonylcarbamoyl group, a lower alkoxybenzenesulfonylcarbamoyl group, a halobenzenesulfonylcarbamoyl group, a mono- or di-lower alkylbenzenesulfonylcarbamoyl group, a C₁-20 alkanesulfonylcarbamoyl group (e.g., 2,2-dimethylethanesulfonylcarbamoyl, butanesulfonylcarbamoyl, propanesulfonylcarbamoyl, isopropanesulfonylcarbamoyl, ethanesulfonylcarbamoyl, methanesulfonylcarbamoyl, octanesulfonylcarbamoyl, pentanesulfonylcarbamoyl, isopentanesulfonylcarbamoyl, hexanesulfonylcarbamoyl, etc.), a trihalo-(lower) alkanesulfonylcarbamoyl group (e.g., trifluoromethanesulfonylcarbamoyl, etc.), a phenyl-(lower) alkanesulfonylcarbamoyl group, a tri-lower alkanesulfonylcarbamoyl group, a lower alkylthio-lower alkanesulfonylcarbamoyl group, a lower alkoxy-(lower) alkanesulfonylcarbamoyl group, a quinolinesulfonylcarbamoyl group, a hydroxy-(lower) alkanesulfonylcarbamoyl group (e.g., 2-hydroxybutanesulfonylcarbamoyl, 3-hydroxybutanesulfonylcarbamoyl, 2-hydroxypentanesulfonylcarbamoyl, etc.), an alkenesulfonylcarbamoyl group (e.g., ethylenesulfonylcarbamoyl, 1-pentenesulfonylcarbamoyl, etc.), a heterocyclic-sulfonylcarbamoyl group (e.g., 2-thiophenesulfonylcarbamoyl, 8-quinolinesulfonylcarbamoyl, etc.).

The lower alkanoyl group is a linear or branched alkylcarbonyl group in which the alkyl moiety has at most 6 carbon atoms. For example, it includes a methylcarbonyl group, an ethylcarbonyl group, an n-propylcarbonyl group, an i-propylcarbonyl group,

an n-butylcarbonyl group, an i-butylcarbonyl group, a sec-butylcarbonyl group, a t-butylcarbonyl group, an n-pentylcarbonyl group, an i-pentylcarbonyl group, a sec-pentylcarbonyl group, a 2,2-dimethylpropylcarbonyl group, a 2-methylbutylcarbonyl group, an n-hexylcarbonyl group, an i-hexylcarbonyl group, a t-hexylcarbonyl group, a sec-hexylcarbonyl group, a 2-methylpentylcarbonyl group, a 3-methylpentylcarbonyl group, a 1-ethylbutylcarbonyl group, a 2-ethylbutylcarbonyl group, a 1,1-dimethylbutylcarbonyl group, a 2,2-dimethylbutylcarbonyl group, a 3,3-dimethylbutylcarbonyl group, a 1-ethyl-1-methylpropylcarbonyl group, etc. Preferred is a carbonyl group bonded to a C₁₋₄ alkyl group, which includes a methylcarbonyl group, an ethylcarbonyl group, an n-propylcarbonyl group, an i-propylcarbonyl group, an n-butylcarbonyl group, an i-butylcarbonyl group, a sec-butylcarbonyl group, a t-butylcarbonyl group, etc.

The aryl group as referred to herein is meant to have from 6 to 10 carbon atoms, including, for example, a phenyl group, a naphthyl group, etc. The naphthyl group includes a 1-naphthyl group, a 2-naphthyl group, etc. On its benzene or naphthalene ring, the aryl group may be substituted by any of a halogen atom, a lower alkyl group, a cyano group, a nitro group, a trifluoromethyl group and the like such as those mentioned above.

The aryl-lower alkyl group is meant to indicate an alkyl group substituted by an aryl group. For the alkyl and aryl moieties in this group, referred to are those mentioned hereinabove. For example, the aryl-lower alkyl group includes a benzyl group, a 1-phenylethyl group, a 2-phenylethyl group, a phenylpropyl group, a phenylbutyl group, a phenylpentyl group, a phenylhexyl group, a naphthylmethyl group, a naphthylethyl group, a naphthylpropyl group, a naphthylbutyl group, a naphthylpentyl group, a naphthylhexyl group, etc.

The aryl-lower alkyloxy group includes, for example, a benzyloxy group, a 1-phenylethyloxy group, a 2-phenylethyloxy group, a phenylpropyloxy group, a phenylbutyloxy group, a phenylpentyloxy group, a phenylhexyloxy group, a naphthylmethyloxy group, a naphthylethyloxy group, a naphthylpropyloxy group, a naphthylbutyloxy group, a naphthylpentyloxy group, etc. On its benzene or naphthalene ring, the aryl-lower alkyloxy group may be substituted.

The aryl-lower alkenyl group is an alkenyl group having at most 6 carbon atoms, to which is bonded an aryl group such as that mentioned above. This includes, for example, a phenylethenyl group, a naphthylethenyl group, etc.

The heterocyclic group is described hereinunder. Concretely, it includes a pyridyl group, a quinolyl group, an isoquinolyl group, a thiazolyl group, a thiadiazolyl group, a benzofuranyl group, a dibenzofuranyl group, a thianaphthalenyl group, a 1H-1,2,3-triazolyl group, a 1,2,4-triazolyl group, a tetrazolyl group, a furyl group, a thienyl group, a pyrrolyl group, an imidazolyl group, a pyrimidinyl group, an indolyl group, a benzimidazolyl group, etc. The heterocyclic group may be optionally substituted by any of a halogen atom, a lower alkyl group or the like mentioned above. For example, the substituted heterocyclic group includes a haloisoquinolyl group, a methylisoquinolyl group, etc.

The pyridyl group as referred to herein is meant to indicate any of a 2-pyridyl group, a 3-pyridyl group or a 4-pyridyl group, in which the bonding position is not specifically defined. The same shall apply to the other heterocyclic groups referred to herein, in which the bonding position is not specifically defined.

Preferably, the "heterocyclic group" indicates a saturated or unsaturated, monocyclic or polycyclic heterocyclic group having at least one hetero atom of oxygen, sulfur and nitrogen atoms and others.

More preferred examples of the heterocyclic group are mentioned below.

A 7- to 12-membered, preferably 9- or 10-membered, unsaturated condensed heterocyclic group (preferably, bicyclic group) having from 1 to 5 nitrogen atoms, for example, an indolyl group, an isoindolyl group, a indolidinyl group, a benzimidazolyl group, a quinolyl group, an isoquinolyl group, an indazolyl group, a benzotriazolyl group, a tetrazolopyridyl group, a tetrazolopyridazinyl group (e.g., tetrazolo[1,5-b]pyridazinyl, etc.), a dihydrotriazolopyridazinyl group, etc.;

A 7- to 12-membered, preferably 9- or 10-membered, unsaturated condensed heterocyclic group (preferably, bicyclic group) having from 1 to 3 sulfur atoms, or its S,S-dioxide, for example, a dithianaphthalenyl group (e.g., 4H-1,3-dithianaphthalenyl, 1,4-dithianaphthalenyl, etc.), a benzothiophenyl group or its S,S-dioxide (e.g., benzo[a]thiophenyl or its S,S-dioxide, benzo[b]thiophenyl or its S,S-dioxide, etc.), etc.;

A 3- to 8-membered, preferably 5- or 6-membered, unsaturated mono-heterocyclic group having from 1 to 4 nitrogen atoms, for example, a pyrrolyl group, a pyrrolinyl group, an imidazolyl group, a pyrazolyl group, a pyridyl group and its N-oxide, a pyrimidyl group, a pyrazinyl group, a pyridazinyl group, a triazolyl group (e.g., 4H-1,2,4-triazolyl, 1H-1,2,3-triazolyl, 2H-1,2,3-triazolyl, etc.), a tetrazolyl group (e.g., 1H-tetrazolyl, 2H-tetrazolyl, etc.), a dihydrotriazinyl group (e.g., 4,5-dihydro-1,2,4-triazinyl, 2,5-dihydro-1,2,4-triazinyl, etc.), etc.;

A 3- to 8-membered, preferably 5- or 6-membered, saturated mono-heterocyclic group having from 1 to 4 nitrogen atoms, for example, an azetidiny group, a pyrrolidinyl group, an imidazolidinyl group, a piperidinyl group, a pyrazolidinyl group, a piperazinyl group, etc.;

A 7- to 12-membered, preferably 9- or 10-membered, unsaturated condensed heterocyclic group (preferably, bicyclic group) having 1 or 2 oxygen atoms and from 1

to 3 nitrogen atoms, for example, a benzoxazolyl group, a benzoxadiazolyl group, etc.;

A 3- to 8-membered, preferably 5- or 6-membered, unsaturated mono-heterocyclic group having 1 or 2 oxygen atoms and from 1 to 3 nitrogen atoms, for example, an oxazolyl group, an isoxazolyl group, an oxadiazolyl group (e.g., 1,2,4-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,5-oxadiazolyl, etc.), etc.;

A 3- to 8-membered, preferably 5- or 6-membered, saturated mono-heterocyclic group having 1 or 2 oxygen atoms and from 1 to 3 nitrogen atoms, for example, a morpholinyl group, etc.;

A 7- to 12-membered, preferably 9- or 10-membered, unsaturated condensed heterocyclic group (preferably, bicyclic group) having 1 or 2 sulfur atoms and from 1 to 3 nitrogen atoms, for example, a benzothiazolyl group, a benzothiadiazolyl group, etc.;

A 3- to 8-membered, preferably 5- or 6-membered, unsaturated mono-heterocyclic group having 1 or 2 sulfur atoms and from 1 to 3 nitrogen atoms, for example, a thiazolyl group, a 1,2-thiazolyl group, a thiazolyl group, a thiadiazolyl group (e.g., 1,2,4-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,5-thiadiazolyl, 1,2,3-thiadiazolyl, etc.), etc.;

A 3- to 8-membered, preferably 5- or 6-membered, saturated mono-heterocyclic group having 1 or 2 sulfur atoms and from 1 to 3 nitrogen atoms, for example, a thiazolidinyl group, etc.;

A 3- to 8-membered, preferably 5- or 6-membered, unsaturated mono-heterocyclic group having one sulfur atom, for example, a thienyl group, etc.

The heterocyclic-lower alkyl group is meant to indicate a lower alkyl group substituted by a heterocyclic group.

The heterocyclic-lower alkyloxy group is meant to indicate an oxy group to which is bonded a heterocyclic-lower alkyl group.

The heterocyclic-lower alkenyl group is meant to indicate a lower alkenyl group substituted by a heterocyclic group.

Preferably, the "acyl group" is an aliphatic acyl group, an aromatic acyl group, a heterocyclic acyl group, or an aromatic group-bonded or heterocyclic group-bonded aliphatic acyl group, which may be derived from carboxylic acids, carbonic acids, sulfonic acids, carbamic acids, etc.

The aliphatic acyl group may be a saturated or unsaturated, acyclic or cyclic one, for example, including an alkanoyl group such as a lower alkanoyl group (e.g., formyl, acetyl, propionyl, butyryl, isobutyryl, valeryl, isovaleryl, pivaloyl, hexanoyl, etc.), an alkylsulfonyl group such as a lower alkylsulfonyl group (e.g., mesyl, ethylsulfonyl, propylsulfonyl, isopropylsulfonyl, butylsulfonyl, isobutylsulfonyl, pentylsulfonyl, hexylsulfonyl, etc.), a carbamoyl group, an N-alkylcarbamoyl group (e.g., methylcarbamoyl, ethylcarbamoyl, etc.), an alkoxycarbonyl group such as a lower alkoxycarbonyl group (e.g., methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, butyloxycarbonyl, tert-butyloxycarbonyl, etc.), an alkenyloxycarbonyl group such as a lower alkenyloxycarbonyl group (e.g., vinyloxycarbonyl, allyloxycarbonyl, etc.), an alkenoyl group such as a lower alkenoyl group (e.g., acryloyl, methacryloyl, crotonoyl, etc.), a cycloalkanecarbonyl group such as a cyclo(lower)-alkanecarbonyl group (e.g., cyclopropanecarbonyl, cyclopentanecarbonyl, cyclohexanecarbonyl, etc.), etc.

The aromatic acyl group includes, for example, a C₆₋₁₀ aroyl group (e.g., benzoyl, toluoyl, xyloyl, etc.), an N-(C₆₋₁₀) arylcarbamoyl group (e.g., N-phenylcarbamoyl, N-tolylcarbamoyl, N-naphthylcarbamoyl, etc.), a C₆₋₁₀ arenesulfonyl group (e.g., benzenesulfonyl, tosyl, etc.), etc.

The heterocyclic acyl group includes, for example, a heterocyclic carbonyl group, a heterocyclic (lower) alkanoyl group (e.g., heterocyclic acetyl, heterocyclic

propanoyl, heterocyclic butanoyl, heterocyclic pentanoyl, heterocyclic hexanoyl, etc.), a heterocyclic (lower) alkenoyl group (e.g., heterocyclic propenoyl, heterocyclic butenoyl, heterocyclic pentenoyl, heterocyclic hexenoyl, etc.), a heterocyclic glyoxyloyl group, a heterocyclic sulfinyl group, a heterocyclic sulfonyl group, etc.

The aromatic group-bonded aliphatic acyl group includes, for example, an aralkoxycarbonyl group such as a phenyl-(lower) alkoxycarbonyl group (e.g., benzyloxycarbonyl, phenethylloxycarbonyl, etc.). etc.

The acyl group may be further substituted by at least one suitable substituent, for example, by a nitro group and the like. As preferred examples of the substituted acyl group, mentioned are a nitroaralkoxycarbonyl group (e.g., nitrobenzyloxycarbonyl, etc.), etc.

Preferred salts of the indole derivatives of the invention are ordinary salts that are not toxic and are acceptable in pharmaceutical use. As their examples, mentioned are salts with bases as well as acid-addition salts, including, for example, salts with inorganic bases such as alkali metal salts with sodium, potassium or the like, alkaline earth metal salts with calcium, magnesium or the like, ammonium salts; organic amine salts with triethylamine, pyridine, picoline, ethanolamine, triethanolamine, dicyclohexylamine, N,N'-dibenzylethyleneamine or the like; inorganic acid salts such as hydrochlorides, hydrobromides, sulfates, phosphates, etc.; organic carboxylates such as formates, acetates, trifluoroacetates, maleates, tartrates, etc.; sulfonic acid-addition salts such as methanesulfonates, benzenesulfonates, p-toluenesulfonates, etc.; salts with basic or acidic amino acids of arginine, aspartic acid, glutamic acid, etc.

The compounds useful in the practice of this invention may have at least one asymmetric center, and therefore they may be in any form of enantiomers or diastereomers. Some alkenyl group-having compounds of the invention may be in any

form of cis- or trans-isomers. In any case, the invention shall encompass their mixtures and individual isomers.

The compounds of the invention may be in any form of tautomers. The invention shall encompass their mixtures and individual tautomers.

The compounds and their salts of the invention may be in any form of their solvates, which shall be within the scope of the invention. Preferably, the solvates are hydrates and ethanolates.

Examples of the compounds of the invention are mentioned below. Specific examples of the indole derivatives of formula (III) include 6-(benzenesulfonylcarbamoyl)-1-(2-chlorobenzyl)-2-methylindole, 1-(biphenyl-4-ylmethyl)-6-(1-butanesulfonylcarbamoyl)-2-ethylindole, 6-(1-butanesulfonylcarbamoyl)-1-(2,4-dichlorobenzyl)-2-methylindole, 1-(2,4-dichlorobenzyl)-2-methyl-6-(1-pentanesulfonylcarbamoyl)indole, 6-(1-butanesulfonylcarbamoyl)-1-(2,4-dichlorobenzyl)-2-ethylindole, 6-(1-butanesulfonylcarbamoyl)-1-(2,4-dichlorobenzyl)indole, 5-(1-butanesulfonylcarbamoyl)-3-(2,4-dichlorobenzyl)indole, 5-(1-butanesulfonylcarbamoyl)-1-(2,4-dichlorobenzyl)indole, 5-(1-butanesulfonylcarbamoyl)-3-(2,4-dichlorobenzyl)-2-methylindole, 5-butanesulfonylcarbamoyl-3-(2,4-dichlorobenzyl)-1-methylindole, 3-(2,4-dichlorobenzyl)-2-methyl-5-(1-pentanesulfonylcarbamoyl)indole, 6-(1-butanesulfonylcarbamoyl)-3-(2,4-dichlorobenzyl)-2-methylindole, 3-(2,4-dichlorobenzyl)-5-(1-pentanesulfonylcarbamoyl)indole, 3-(2,4-dichlorobenzyl)-2-methyl-5-(1-propanesulfonylcarbamoyl)indole, 3-(2,4-dichlorobenzyl)-2-methyl-5-(1-octanesulfonylcarbamoyl)indole, 5-(benzenesulfonylcarbamoyl)-3-(2,4-dichlorobenzyl)-2-methylindole, 3-(2,4-dichlorobenzyl)-5-(1-

[illegible]

dichlorobenzyl)-2-methylindole, 3-(2,4-dichlorobenzyl)-2-methyl-5-(propanesultam-1-ylcarbonyl)indole, 6-(1-butanesulfonylcarbamoyle)-2-(2,4-dichlorobenzyl)-3-methylindole, 1-(2,4-dichlorobenzyl)-3-methyl-6-(1-pentanesulfonylcarbamoyle)indole, 3-(2,4-dichlorobenzyl)-2-methyl-5-((4-methylbenzene)sulfonylcarbamoyle)indole, 3-(2,4-dichlorobenzyl)-2-methyl-5-((4-nitrobenzene)sulfonylcarbamoyle)indole, 5-((4-chlorobenzene)sulfonylcarbamoyle)-3-(2,4-dichlorobenzyl)-2-methylindole, 5-((3-chlorobenzene)sulfonylcarbamoyle)-3-(2,4-dichlorobenzyl)-2-methylindole, 5-((2-chlorobenzene)sulfonylcarbamoyle)-3-(2,4-dichlorobenzyl)-2-methylindole, 3-(2,4-dichlorobenzyl)-5-((4-fluorobenzene)sulfonylcarbamoyle)-2-methylindole, 3-(2,4-dichlorobenzyl)-2-methyl-5-((2-naphthalene)sulfonylcarbamoyle)indole, 3-(2,4-dichlorobenzyl)-2-methyl-5-((1-naphthalene)sulfonylcarbamoyle)indole, 3-(2,4-dichlorobenzyl)-2-methyl-5-((2-methylbenzene)sulfonylcarbamoyle)indole, 3-(2,4-dichlorobenzyl)-5-(2,6-dimethylbenzene)sulfonylcarbamoyle-2-methylindole, 5-(4-bromobenzene)sulfonylcarbamoyle-3-(2,4-dichlorobenzyl)-2-methylindole, 3-(2,4-dichlorobenzyl)-2-methyl-5-((E)-styrenesulfonylcarbamoyle)indole, 3-(2,4-dichlorobenzyl)-2-methyl-5-((4-vinylbenzene)sulfonylcarbamoyle)indole, 5-((4-phenylazobenzene)sulfonylcarbamoyle)-3-(2,4-dichlorobenzyl)-2-methylindole, 3-(2,4-dichlorobenzyl)-2-methyl-5-((4-trifluoromethylbenzene)sulfonylcarbamoyle)indole, 3-(2,4-dichlorobenzyl)-2-methyl-5-((4-methyl-1-pent-1-ene)sulfonylcarbamoyle)indole, 3-(2,4-dichlorobenzyl)-5-((3,4-dimethoxybenzene)sulfonylcarbamoyle)-2-methylindole, 5-((4-t-butylbenzene)sulfonylcarbamoyle)-3-(2,4-dichlorobenzyl)-2-methylindole, 3-(2,4-dichlorobenzyl)-2-methyl-5-((3-methylbenzene)sulfonylcarbamoyle)indole, 3-(2,4-dichlorobenzyl)-2-methyl-5-(2-octanesulfonylcarbamoyle)indole, 3-(2,4-dichlorobenzyl)-2-methyl-5-((4-phenylbenzene)sulfonylcarbamoyle)indole, 3-((2-chloro-4-phenyl)benzyl)-2-methyl-5-((1-pent-1-ene)sulfonylcarbamoyle)indole, 3-(2-chloro-4-

phenylbenzyl)-2-methyl-5-(1-pentanesulfonylcarbamoyle)indole, 5-
 (benzenesulfonylcarbamoyle)-3-((2-chloro-4-phenyl)benzyl)-2-methylindole, 3-(2,4-
 dichlorobenzyl)-5-((4-ethylbenzene)sulfonylcarbamoyle)-2-methylindole, 5-((4-n-
 butylbenzene)sulfonylcarbamoyle)-3-(2,4-dichlorobenzyl)-2-methylindole, 5-((4-n-
 butoxybenzene)sulfonylcarbamoyle)-3-(2,4-dichlorobenzyl)-2-methylindole, 3-(2,4-
 dichlorobenzyl)-2-methylthio-5-((1-pent-1-ene)sulfonylcarbamoyle)indole, 5-
 (benzenesulfonylcarbamoyle)-3-(2,4-dichlorobenzyl)-2-methylthioindole, 3-(2,4-
 dichlorobenzyl)-2-methylthio-5-(1-pentanesulfonylcarbamoyle)indole, 3-(2,4-
 dichlorobenzyl)-2-methyl-5-((1-penta-1,3-diene)sulfonylcarbamoyle)indole, 5-((2-
 cyclopropylethylene)sulfonylcarbamoyle)-3-(2,4-dichlorobenzyl)-2-methylindole, 3-
 (2,4-dichlorobenzyl)-2-methyl-5-((4-methyl-(E)-styrene)sulfonylcarbamoyle)indole,
 3-(2,4-dichlorobenzyl)-5-((4-methoxy-(E)-styrene)sulfonylcarbamoyle)-2-
 methylindole, 3-(2,4-dichlorobenzyl)-2-methoxymethyl-5-(1-
 pentanesulfonylcarbamoyle)indole, 3-((1-bromonaphthalen-2-yl)methyl)-2-methyl-5-
 ((E)-styrene)sulfonylcarbamoyle)indole, 3-((1-bromonaphthalen-2-yl)methyl)-2-
 methyl-5-(4-vinylbenzene)sulfonylcarbamoyle)indole, 3-((1-bromonaphthalen-2-
 yl)methyl)-2-methyl-5-(p-toluenesulfonylcarbamoyle)indole, 5-
 (benzenesulfonylcarbamoyle)-3-((1-bromonaphthalen-2-yl)methyl)-2-methylindole, 3-
 ((2-chloro-4-phenyl)benzyl)-2-methyl-5-((E)-styrenesulfonylcarbamoyle)indole, 3-
 ((2-chloro-4-phenyl)benzyl)-2-methyl-5-((4-vinylbenzene)sulfonylcarbamoyle)indole, 3-
 ((1-bromonaphthalen-2-yl)methyl)-2-methyl-5-((1-pent-1-
 ene)sulfonylcarbamoyle)indole, 3-((2-chloro-4-phenyl)benzyl)-2-methyl-5-(p-
 toluenesulfonylcarbamoyle)indole, 3-(4-bromo-2-chlorobenzyl)-2-methyl-5-(1-
 pentanesulfonylcarbamoyle)indole, 3-(4-bromo-2-chlorobenzyl)-2-methyl-5-(2-(5-
 chlorothienyl)sulfonylcarbamoyle)indole, 3-(2-chloro-4-nitrobenzyl)-2-methyl-5-(1-

pentanesulfonylcarbamoyl)indole, 3-(2-chloro-4-(2-phenylethenyl)benzyl)-2-methyl-5-(1-pentanesulfonylcarbamoyl)indole, etc.

Specific examples of the indole derivatives of formula (V) include 1-(2-chlorobenzyl)-6-methoxycarbonyl-2-methylindole, 6-carboxy-1-(2-chlorobenzyl)-2-methylindole, 1-(biphenyl-4-ylmethyl)-2-ethyl-6-methoxycarbonylindole, 1-(biphenyl-4-ylmethyl)-6-carboxy-2-ethylindole, 1-(2,4-dichlorobenzyl)-6-methoxycarbonyl-2-methylindole, 6-carboxy-1-(2,4-dichlorobenzyl)-2-methylindole, 1-(2,4-dichlorobenzyl)-2-ethyl-6-methoxycarbonylindole, 6-carboxy-1-(2,4-dichlorobenzyl)-2-ethylindole, 1-(2,4-dichlorobenzyl)-6-methoxycarbonylindole, 6-carboxy-1-(2,4-dichlorobenzyl)indole, 3-(2,4-dichlorobenzyl)-5-methoxycarbonylindole, 5-carboxy-3-(2,4-dichlorobenzyl)indole, 1-(2,4-dichlorobenzyl)-5-methoxycarbonylindole, 5-carboxy-1-(2,4-dichlorobenzyl)indole, 3-(2,4-dichlorobenzyl)-5-methoxycarbonyl-2-methylindole, 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole, 3-(2,4-dichlorobenzyl)-5-methoxycarbonyl-1-methylindole, 5-carboxy-3-(2,4-dichlorobenzyl)-1-methylindole, 3-(2,4-dichlorobenzyl)-6-(methoxycarbonyl)-2-methylindole, 3-(biphenyl-3-ylmethyl)-5-(methoxycarbonyl)-2-methylindole, 3-(2-chlorobenzyl)-5-(methoxycarbonyl)-2-methylindole, 3-(2,4-dichlorobenzyl)-5-(methoxycarbonyl)-2-methylindole, 3-(4-benzyloxybenzyl)-5-(methoxycarbonyl)-2-methylindole, 3-(2,4-dichlorobenzyl)-5-(methoxycarbonyl)-2-propylindole, 3-(2,4-dichlorobenzyl)-2-ethyl-5-(methoxycarbonyl)indole, 3-(2,4-dichlorobenzyl)-5-(methoxycarbonyl)indole, 3-(1-bromonaphthalen-2-ylmethyl)-5-(methoxycarbonyl)-2-methylindole, 3-((3-chloropyridin-4-yl)methyl)-5-(methoxycarbonyl)-2-methylindole, 5-(methoxycarbonyl)-2-methyl-3-(4-(2-phenylethenyl)benzyl)indole, 3-((4-chloroisoquinolin-3-yl)methyl)-5-(methoxycarbonyl)-2-methylindole, 3-((4-bromoisoquinolin-3-yl)methyl)-5-(methoxycarbonyl)-2-methylindole, 2-(2,4-

dichlorobenzoyl)-6-(methoxycarbonyl)-3-methylindole, 1-(2,4-dichlorobenzyl)-6-(methoxycarbonyl)-3-methylindole, 6-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole, 3-(biphenyl-4-ylmethyl)-5-carboxy-2-methylindole, 5-carboxy-3-(2-chlorobenzyl)-2-methylindole, 5-carboxy-3-(2,4-dichlorobenzoyl)-2-methylindole, 3-(4-benzyloxybenzyl)-5-carboxy-2-methylindole, 5-carboxy-3-(2,4-dichlorobenzyl)-2-propylindole, 5-carboxy-3-(2,4-dichlorobenzyl)-2-ethylindole, 5-carboxy-3-(2,4-dichlorobenzyl)indole, 3-(1-bromonaphthalen-2-ylmethyl)-5-carboxy-2-methylindole, 5-carboxy-3-((3-chloropyridin-4-yl)methyl)-2-methylindole, 5-carboxy-2-methyl-3-(4-(2-phenylethenyl)benzyl)indole, 5-carboxy-3-((4-chloroisoquinolin-3-yl)methyl)-2-methylindole, 3-((4-bromoisoquinolin-3-yl)methyl)-5-carboxy-2-methylindole, 6-carboxy-2-(2,4-dichlorobenzyl)-3-methylindole, 6-carboxy-1-(2,4-dichlorobenzyl)-3-methylindole, 3-(2-chloro-4-phenylbenzyl)-5-methoxycarbonyl-2-methylindole, 5-carboxy-3-(2-chloro-4-phenylbenzyl)-2-methylindole, 3-(2,4-dichlorobenzyl)-5-methoxycarbonyl-2-methylthioindole, 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylthioindole, 3-(2,4-dichlorobenzyl)-5-methoxycarbonyl-2-(methoxymethyl)indole, 5-carboxy-3-(2,4-dichlorobenzyl)-2-(methoxymethyl)indole, etc.

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 shows the chemical formulae of compounds (30) to (34).

Fig. 2 shows the chemical formulae of compounds (35) to (39).

Fig. 3 shows the chemical formulae of compounds (40) to (44).

Fig. 4 shows the chemical formulae of compounds (45) to (49).

Fig. 5 shows the chemical formulae of compounds (50) to (54).

Fig. 6 shows the chemical formulae of compounds (55) to (59).

Fig. 7 shows the chemical formulae of compounds (60) to (64).

Fig. 8 shows the chemical formulae of compounds (65) to (69).

Fig. 9 shows the chemical formulae of compounds (70) to (74).

Fig. 10 shows the chemical formulae of compounds (75) to (79).

Fig. 11 shows the chemical formulae of compounds (80) to (84).

Fig. 12 shows the chemical formulae of compounds (85) to (89).

Fig. 13 shows the chemical formulae of compounds (90) to (94).

Fig. 14 shows the chemical formulae of compounds (95) to (99).

Fig. 15 shows the chemical formulae of compounds (100) to (104).

Fig. 16 shows the chemical formulae of compounds (105) to (109).

Fig. 17 shows the chemical formulae of compounds (110) to (114).

Fig. 18 shows the chemical formulae of compounds (115) to (119).

Fig. 19 shows the chemical formulae of compounds (120) to (124).

Fig. 20 shows the chemical formulae of compounds (125) to (129).

Fig. 21 shows the chemical formulae of compounds (130) to (134).

Fig. 22 shows the chemical formulae of compounds (135) to (139).

Fig. 23 shows the chemical formulae of compounds (140) to (144).

Fig. 24 shows the chemical formulae of compounds (145) to (149).

Fig. 25 shows the chemical formulae of compounds (150) to (154).

Fig. 26 shows the chemical formulae of compounds (155) to (159).

Fig. 27 shows the chemical formulae of compounds (160) to (164).

Fig. 28 shows the chemical formulae of compounds (165) to (169).

Fig. 29 shows the chemical formulae of compounds (170) to (173).

The synthesis of the compounds useful in this invention and of the intermediates for use in making those compounds are illustrated by the following, non-limiting Examples from the above PCT Application WO 98/15530.

Production Example 1; Methyl 4-(2-Oxopropyl)-3-Nitrobenzoate

To a methylene chloride (60 ml) solution of methyl E-4-(2-dimethylaminovinyl)-3-nitrobenzoate (5.0 g) (this is reportedly produced according to the method of US-5212195) and pyridine (2.6 ml), added are acetyl chloride (2 ml), and stirred at room temperature for 20 hours. Water is added to this to stop the reaction, and the organic layer is concentrated. To the residue, added is 1,4-dioxane (25 ml) and water (13 ml), and heated under reflux for 16 hours. The solvent is evaporated away under reduced pressure, and the residue is extracted with ethyl acetate and water. The organic layer is washed with a saturated saline solution, then concentrated, and purified through silica gel column chromatography (eluent: hexane/ethyl acetate = 2/1) to obtain methyl 4-(2-oxopropyl)-3-nitrobenzoate (4.12 g).

$^1\text{H-NMR}$ (CDCl_3 , δ): 2.35 (3H, s), 3.97 (3H, s), 4.19 (2H, s), 7.37 (1H, d, $J = 7.9$ Hz), 8.24 (1H, d, $J = 7.9$ Hz), 8.75 (1H, s).

Production Example 2; 6-Methoxycarbonyl-2-Methylindole

A mixed solution of methyl 4-(2-oxopropyl)-3-nitrobenzoate (3.86 g), reduced iron (9.0 g) and acetic acid (40 ml) is stirred at 90°C for 24 hours. The solid is removed through filtration, and the filtrate is concentrated under reduced pressure. The resulting residue is purified through silica gel column chromatography (eluent: hexane/ethyl acetate = 3/1) to obtain 6-methoxycarbonyl-2-methylindole (0.890 g).

[Physical Properties of the Product]

$^1\text{H-NMR}$ (CDCl_3 , δ): 2.46 (3H, s), 3.90 (3H, s), 6.25 (1H, s), 7.48 (1H, d, $J = 8.3$ Hz), 7.72 (1H, dd, $J = 1.4$ Hz and 8.3 Hz), 7.97 (1H, s), 8.39 (1H, brs).

<Example 1; Production of 1-(2-chlorobenzyl)-6-methoxycarbonyl-2-methylindole (30)>

A mixture of 6-methoxycarbonyl-2-methylindole (0.89 g), 2-chlorobenzyl

bromide (1.45 g), potassium carbonate (0.780 g) and N,N-dimethylformamide (2 ml) is stirred at 80°C for 18 hours. The solvent is evaporated away under reduced pressure, and the resulting residue is purified through silica gel column chromatography (eluent: hexane/chloroform = 3/1) to obtain 1-(2-chlorobenzyl)-6-methoxycarbonyl-2-methylindole (30) (0.660 g).

[Physical Properties of Compound (30)]

¹H-NMR (CDCl₃, δ): 2.35 (3H, s), 3.88 (3H, s), 5.41 (2H, s), 6.17 (1H, d, J = 7.8 Hz), 6.42 (1H, s), 7.02 (1H, t, J = 7.6 Hz), 7.18 (1H, t, J = 7.1 Hz), 7.42 (1H, d, J = 8.0 Hz), 7.58 (1H, d, J = 8.3 Hz), 7.80 (1H, dd, J = 1.2 and 8.3 Hz).

<Example 2; Production of 6-carboxy-1-(2-chlorobenzyl)-2-methylindole (31)>

A mixture of 1-(2-chlorobenzyl)-6-methoxycarbonyl-2-methylindole (0.630 g), aqueous 10 % sodium hydroxide solution (2.7 g), water (1.8 g) and ethanol (7.0 g) is refluxed for 2.5 hours. After cooled, this is processed with 10 % HCl to have a pH of 4, and the crystal formed is taken out through filtration and dried to obtain 6-carboxy-1-(2-chlorobenzyl)-2-methylindole (31) (0.480 g).

[Physical Properties of Compound (31)]

¹H-NMR (DMSO-d₆, δ): 2.35 (3H, s), 5.53 (2H, s), 6.13 (1H, d, J = 7.6 Hz), 6.46 (1H, s), 7.15 (1H, t, J = 7.6 Hz), 7.28 (1H, t, J = 7.6 Hz), 7.54 (2H, t, J = 8.3 Hz), 7.62 (1H, d, J = 8.3 Hz), 12.49 (1H, brs).

<Example 3; Production of 6-benzenesulfonylcarbonyl-1-(2-chlorobenzyl)-2-methylindole (32)>

To an N,N-dimethylformamide (10 ml) solution of 6-carboxy-1-(2-chlorobenzyl)-2-methylindole (0.470 g), added is N,N'-carbonyldiimidazole (0.508 g), and stirred at room temperature for 1 hour. Next, benzenesulfonamide (0.493 g) and diazabicycloundecene (0.477 g) are added thereto, and stirred at 100°C for 72 hours.

The solvent is evaporated away under reduced pressure, and the residue is purified through silica gel column chromatography (eluent: chloroform) to obtain 6-(benzenesulfonylcarbamoyl)-1-(2-chlorobenzyl)-2-methylindole. This is dissolved in a small amount of ethyl acetate, to which is added hexane. The crystal formed is taken out through filtration, and dried to obtain 6-(benzenesulfonylcarbamoyl)-1-(2-chlorobenzyl)-2-methylindole (32) (0.440 g).

[Physical Properties of Compound (32)]

$^1\text{H-NMR}$ (DMSO- d_6 , δ): 2.30 (3H, s), 5.50 (2H, s), 6.05 (1H, d, $J = 7.7$ Hz), 6.45 (1H, s), 7.14 (1H, t, $J = 7.6$ Hz), 7.29 (1H, t, $J = 7.7$ Hz), 7.48 - 7.68 (6H, m), 7.89 - 8.01 (3H, m), 12.22 (1H, brs).

<Production Example 3; methyl 4-(2-oxobutyl)-3-nitrobenzoate>

According to the method of Production Example 1, a mixed solution of methyl E-4-(2-dimethylaminovinyl)-3-nitrobenzoate (6.30 g), pyridine (3.2 ml), propionyl chloride (3.26 g) and methylene chloride (60 ml) is heated under reflux for 11 hours, and then heated under reflux with 1,4-dioxane in water for 16 hours to obtain methyl 4-(2-oxobutyl)-3-nitrobenzoate (5.63 g).

[Physical Properties of the Product]

$^1\text{H-NMR}$ (CDCl_3 , δ): 1.12 (3H, t, $J = 7.3$ Hz), 2.65 (2H, q, $J = 7.3$ Hz), 3.97 (3H, s), 4.18 (2H, s), 7.37 (1H, d, $J = 7.9$ Hz), 8.22 (1H, dd, $J = 1.7$ and 7.9 Hz), 8.73 (1H, d, $J = 1.7$ Hz).

<Production Example 4; 2-ethyl-6-methoxycarbonylindole>

According to the method of Production Example 2, obtained is 2-ethyl-6-methoxycarbonylindole (2.55 g) from methyl 4-(2-oxobutyl)-3-nitrobenzoate (5.60 g) and reduced iron (13.0 g).

[Physical Properties of the Product]

$^1\text{H-NMR}$ (CDCl_3 , δ): 1.37 (3H, t, $J = 7.6$ Hz), 2.83 (2H, q, $J = 7.6$ Hz), 3.92 (3H, s), 6.30 (1H, s), 7.53 (1H, d, $J = 8.3$ Hz), 7.76 (1H, d, $J = 8.3$ Hz), 8.05 (1H, s), 8.20 (1H, brs).

<Example 4: Production of 1-(biphenyl-4-ylmethyl)-2-ethyl-6-methoxycarbonylindole (33)>

According to the method of Example 1, obtained is 1-(biphenyl-4-ylmethyl)-2-ethyl-6-methoxycarbonylindole (33) (2.50 g) from 2-ethyl-6-methoxycarbonylindole (2.55 g), 4-biphenylmethyl bromide (4.00 g) and potassium carbonate (2.07 g). This is directly used in the next reaction.

<Example 5; Production of 1-(biphenyl-4-ylmethyl)-6-carboxy-2-ethylindole (34)>

According to the method of Example 3, obtained is 1-(biphenyl-4-ylmethyl)-6-carboxy-2-ethylindole (34) (1.29 g) from 1-(biphenyl-4-ylmethyl)-2-ethyl-6-methoxycarbonylindole (2.50 g).

[Physical Properties of Compound (34)]

$^1\text{H-NMR}$ (DMSO-d_6 , δ): 1.26 (3H, t, $J = 7.5$ Hz), 2.76 (2H, q, $J = 7.5$ Hz), 5.30 (2H, s), 6.24 (1H, s), 7.35 - 7.74 (11H, m), 7.94 (1H, s), 12.57 (1H, brs).

<Example 6; Production of 1-(biphenyl-4-ylmethyl)-6-(1-butanefulfonylcarbonyl)-2-ethylindole (35)>

According to the method of Example 3, obtained is 1-(biphenyl-4-ylmethyl)-6-(1-butanefulfonylcarbonyl)-2-ethylindole (35) (0.360 g) from 1-(biphenyl-4-ylmethyl)-6-carboxy-2-ethylindole (0.400 g), N,N' -carbonyldiimidazole (0.365 g), 1-butanefulfonylamine (0.309 g) and diazabicycloundecene (0.343 g).

[Physical Properties of Compound (35)]

$^1\text{H-NMR}$ (DMSO-d_6 , δ): 0.86 (3H, t, $J = 7.4$ Hz), 1.23 (3H, t, $J = 7.5$ Hz), 1.37 (2H, m), 1.64 (2H, m), 2.70 (2H, q, $J = 7.4$ Hz), 3.28 (2H, m), 5.31 (2H, s), 6.17

(1H, s), 7.39 (1H, t, J = 7.4 Hz), 7.43 - 7.51 (3H, m), 7.59 (2H, d, J = 8.2 Hz), 7.66 - 7.72 (3H, m), 7.75 (2H, d, J = 8.2 Hz), 8.11 (1H, d), 11.96 (1H, brs).

IR (Nujol): 1650 cm^{-1} .

m.p.: 71 - 83°C.

<Example 7; Production of 1-(2,4-dichlorobenzyl)-6-methoxycarbonyl-2-methylindole (36)>

According to the method of Example 1, obtained is 1-(2,4-dichlorobenzyl)-6-methoxycarbonyl-2-methylindole (36) (1.71 g) from 6-methoxycarbonyl-2-methylindole (1.59 g), 2,4-dichlorobenzyl chloride (2.14 g), potassium carbonate (1.51 g) and sodium iodide (1.26 g).

$^1\text{H-NMR}$ (CDCl_3 , δ): 2.31 (3H, s), 3.94 (3H, s), 5.29 (2H, s), 6.11 (1H, s), 7.26 (1H, dd, J = 2.0 and 8.2 Hz), 7.32 (1H, d, J = 8.1 Hz), 7.48 (2H, m), 7.76 (1H, d, J = 8.3 Hz), 8.02 (1H, s).

<Example 8; Production of 6-carboxy-1-(2,4-dichlorobenzyl)-2-methylindole (37)>

According to the method of Example 1, obtained is 6-carboxy-1-(2,4-dichlorobenzyl)-2-methylindole (37) (1.21 g) from 1-(2,4-dichlorobenzyl)-6-methoxycarbonyl-2-methylindole (1.70 g).

$^1\text{H-NMR}$ (DMSO-d_6 , δ): 2.35 (3H, s), 5.35 (2H, s), 6.22 (1H, s), 7.45 - 7.50 (2H, m), 7.55 - 7.61 (2H, m), 7.74 (1H, d, J = 2.1 Hz), 7.87 (1H, s), 12.57 (1H, brs).

<Example 9; Production of 6-(1-butanefulfonylcarbonyl)-1-(2,4-dichlorobenzyl)-2-methylindole (38)>

According to the method of Example 3, obtained is 1-(2,4-dichlorobenzyl)-6-(1-butanefulfonylcarbonyl)-2-methylindole (38) (0.270 g) from 6-carboxy-1-(2,4-dichlorobenzyl)-2-methylindole (0.370 g), N,N'-carbonyldiimidazole (0.269 g), 1-

butanesulfonamide (0.228 g) and diazabicycloundecene (0.253 g).

¹H-NMR (CDCl₃, δ): 0.90 (3H, t), 1.44 (2H, m), 1.82 (2H, m), 3.10 (2H, m), 5.26 (2H, s), 6.11 (1H, s), 7.20 - 7.30 (2H, m), 7.42 - 7.49 (2H, m), 7.52 - 7.59 (1H, m), 7.97 (1H, s).

IR (neat): 1682 cm⁻¹.

Oily.

<Example 10; Production of 1-(2,4-dichlorobenzyl)-2-methyl-6-(1-pentanesulfonylcarbamoyl)indole (39)>

According to the method of Example 3, obtained is 1-(2,4-dichlorobenzyl)-2-methyl-6-(1-pentanesulfonylcarbamoyl)indole (39) (0.482 g) from 6-carboxy-1-(2,4-dichlorobenzyl)-2-methylindole (0.668 g), N,N'-carbonyldiimidazole (0.649 g), 1-pentanesulfonamide (0.605 g) and diazabicycloundecene (0.609 g).

¹H-NMR (DMSO-d₆, δ): 0.85 (3H, t, J = 7.3 Hz), 1.30 (2H, m), 1.39 (2H, m), 1.72 (2H, m), 2.28 (3H, s), 3.53 (2H, t, J = 7.7 Hz), 5.43 (2H, s), 6.23 (1H, s), 7.51 (1H, dd, J = 2.0 and 8.2 Hz), 7.53 (1H, d, J = 8.4 Hz), 7.59 (1H, d, J = 8.4 Hz), 7.61 (1H, dd, J = 2.0 and 8.2 Hz), 7.77 (1H, d, J = 2.0 Hz), 8.04 (1H, s), 11.94 (1H, s).

IR (neat): 1685 cm⁻¹.

Oily.

<Example 11; Production of 1-(2,4-dichlorobenzyl)-2-ethyl-6-methoxycarbonylindole (40)>

According to the method of Example 1, obtained is 1-(2,4-dichlorobenzyl)-2-ethyl-6-methoxycarbonylindole (40) (2.70 g) from 2-ethyl-6-methoxycarbonylindole (4.88 g), 2,4-dichlorobenzyl chloride (6.10 g), potassium carbonate (4.31 g) and potassium iodide (3.98 g).

$^1\text{H-NMR}$ (CDCl_3 , δ): 1.33 (3H, t, $J = 7.4$ Hz), 2.64 (2H, t, $J = 7.4$ Hz), 3.89 (3H, s), 5.37 (2H, s), 6.08 (1H, d, $J = 8.4$ Hz), 6.45 (1H, s), 6.99 (1H, dd, $J = 2.0$ and 8.4 Hz), 7.45 (1H, d, $J = 2.0$ Hz), 7.61 (1H, d, $J = 8.3$ Hz), 7.81 (1H, d, $J = 8.3$ Hz), 7.88 (1H, s).

<Example 12; Production of 6-carboxy-1-(2,4-dichlorobenzyl)-2-ethylindole (41)>

According to the method of Example 2, obtained is 6-carboxy-1-(2,4-dichlorobenzyl)-2-ethylindole (41) (0.837 g) from 1-(2,4-dichlorobenzyl)-2-ethyl-6-methoxycarbonylindole (0.959 g).

$^1\text{H-NMR}$ (DMSO-d_6 , δ): 1.24 (3H, t, $J = 7.4$ Hz), 2.67 (2H, q, $J = 7.5$ Hz), 5.53 (2H, s), 6.09 (1H, d, $J = 8.4$ Hz), 6.48 (1H, s), 7.26 (1H, d, $J = 8.4$ Hz), 7.59 (1H, d, $J = 8.3$ Hz), 7.64 (1H, d, $J = 8.3$ Hz), 7.72 (1H, d, $J = 2.0$ Hz), 7.88 (1H, s).

<Example 13: Production of 6-(1-butanefonylcarbonyl)-1-(2,4-dichlorobenzyl)-2-ethylindole (42)>

According to the method of Example 3, obtained is 6-(1-butanefonylcarbonyl)-1-(2,4-dichlorobenzyl)-2-ethylindole (42) (0.220 g) from 6-carboxy-1-(2,4-dichlorobenzyl)-2-ethylindole (0.801 g), N,N' -carbonyldiimidazole (0.567 g), 1-butanefonyl (0.480 g) and diazabicycloundecene (0.533 g).

$^1\text{H-NMR}$ (DMSO-d_6 , δ): 0.84 (3H, t, $J = 7.3$ Hz), 1.22 (3H, t, $J = 7.4$ Hz), 1.38 (2H, m), 1.64 (2H, m), 2.63 (2H, m), 3.48 (2H, m), 5.51 (2H, s), 6.05 (1H, d, $J = 8.5$ Hz), 6.50 (1H, s), 7.26 (1H, dd, $J = 1.9$ and 8.4 Hz), 7.63 (2H, m), 7.72 (1H, d, $J = 2.0$ Hz), 8.06 (1H, s), 11.70 (1H, s).

IR (Nujol): 1682 cm^{-1} .

m.p.: $162.2 - 162.7^\circ\text{C}$.

<Production Example 5; 6-methoxycarbonylindole>

To a mixed solvent of tetrahydrofuran (30 ml), ethanol (30 ml) and water (100

ml), added are methyl E-4-(2-dimethylaminovinyl)-3-nitrobenzoate (10.0 g) and sodium hydrosulfite (104.5 g), and stirred at 70°C for 1 hours. After this is cooled to room temperature, a saturated saline solution is added thereto. Then, this is extracted with chloroform. The organic layer is dried, and the solvent is evaporated away. The resulting residue is purified through silica gel column chromatography (eluent: hexane/ethyl acetate = 2/1 to 1/1) to obtain 6-methoxycarbonylindole (2.79 g).

$^1\text{H-NMR}$ (CDCl_3 , δ): 3.93 (3H, s), 6.60 (1H, s), 7.37 (1H, m), 7.66 (1H, d, $J = 8.3$ Hz), 7.81 (1H, dd, $J = 1.3$ and 8.3 Hz), 8.17 (1H, s), 8.52 (1H, brs).

<Example 14; Production of 1-(2,4-dichlorobenzyl)-6-methoxycarbonylindole (43)>

According to the method of Example 1, obtained is 1-(2,4-dichlorobenzyl)-6-methoxycarbonylindole (43) (1.97 g) from 6-methoxycarbonylindole (2.79 g), 2,4-dichlorobenzyl chloride (4.67 g), potassium carbonate (3.30 g) and potassium iodide (3.96 g).

$^1\text{H-NMR}$ (CDCl_3 , δ): 3.91 (3H, s), 5.43 (2H, s), 6.47 (1H, d, $J = 8.3$ Hz), 6.63 (1H, d, $J = 3.2$ Hz), 7.07 (1H, dd, $J = 1.3$ and 8.4 Hz), 7.26 (1H, d, $J = 1.0$ Hz), 7.45 (1H, d, $J = 1.5$ Hz), 7.68 (1H, d, $J = 8.3$ Hz), 7.83 (1H, dt, $J = 1.0$ and 8.4 Hz), 8.01 (1H, s).

<Example 15; Production of 6-carboxy-1-(2,4-dichlorobenzyl)indole (44)>

According to the method of Example 2, obtained is 6-carboxy-1-(2,4-dichlorobenzyl)indole (44) (1.55 g) from 1-(2,4-dichlorobenzyl)-6-methoxycarbonylindole (1.67 g).

[Physical Properties of Compound (44)]

$^1\text{H-NMR}$ (DMSO-d_6 , δ): 5.59 (2H, s), 6.61 (2H, m), 7.32 (1H, d, $J = 8.2$ Hz), 7.65 (3H, s), 7.69 (1H, s), 8.00 (1H, s).

<Example 16; Production of 6-(1-butanesulfonylcarbamoyl)-1-(2,4-

dichlorobenzyl)indole (45)>

According to the method of Example 3, obtained is 6-(1-butanefonylcarbonyl)-1-(2,4-dichlorobenzyl)indole (45) (0.307 g) from 6-carboxy-1-(2,4-dichlorobenzyl)indole (0.800 g), N,N'-carbonyldiimidazole (0.567 g), 1-butanefonylamine (0.480 g) and diazabicycloundecene (0.533 g).

$^1\text{H-NMR}$ (DMSO- d_6 , δ): 0.84 (3H, m), 1.38 (2H, m), 1.65 (2H, m), 3.47 (2H, m), 5.57 (2H, s), 6.53 (1H, d, $J = 8.4$ Hz), 6.64 (1H, d, $J = 3.0$ Hz), 7.32 (1H, dd, $J = 2.0$ and 8.4 Hz), 7.66 (3H, m), 7.71 (1H, d, $J = 2.0$ Hz), 8.13 (1H, s), 11.79 (1H, s).

IR (neat): 1694 cm^{-1} .

Oily.

<Production Example 6; 2,4-dichlorobenzyl iodide>

2,4-Dichlorobenzyl chloride (23.45 g) and potassium iodide (47.88 g) are stirred in acetone (100 ml) at room temperature for 24 hours. After concentrated, this is extracted with *t*-butyl methyl ether (300 ml) and water (100 ml). The extract is washed with water, dried and concentrated to obtain 2,4-dichlorobenzyl iodide (34.49 g).

$^1\text{H-NMR}$ (CDCl_3 , δ): 4.48 (2H, s), 7.19 (1H, dd, $J = 2.2$ and 8.3 Hz), 7.34 (1H, d, $J = 8.3$ Hz), 7.36 (1H, d, $J = 2.2$ Hz).

<Example 17; Production of 3-(2,4-dichlorobenzyl)-5-methoxycarbonylindole (46)>

Silver oxide (28.09 g) is added to a dioxane (200 ml) solution of 5-methoxycarbonylindole (17.52 g) and 2,4-dichlorobenzyl iodide (34.49 g), and stirred under reflux for 6.5 hours. The insoluble is removed through filtration, and the filtrate is concentrated under reduced pressure. The resulting oily residue is purified through silica gel column chromatography (eluent: hexane/ethyl acetate = 4/1), and recrystallized from hexane/ethyl acetate to obtain 3-(2,4-dichlorobenzyl)-5-

methoxycarbonylindole (46) (5.49 g).

$^1\text{H-NMR}$ (CDCl_3 , δ): 3.92 (3H, s), 4.20 (2H, s), 6.98 (1H, d, $J = 2.3$ Hz), 7.09 - 7.13 (2H, m), 7.38 (1H, d, $J = 9.1$ Hz), 7.42 (1H, d, $J = 1.9$ Hz), 7.92 (1H, dd, $J = 1.5$ and 8.5 Hz), 8.22 (1H, brs), 8.31 (1H, d, $J = 1.2$ Hz).

<Example 18; Production of 5-carboxy-3-(2,4-dichlorobenzyl)indole (47)>

According to the method of Example 2, obtained is 5-carboxy-3-(2,4-dichlorobenzyl)indole (47) (4.64 g) from 3-(2,4-dichlorobenzyl)-5-methoxycarbonylindole (5.02 g).

$^1\text{H-NMR}$ (DMSO-d_6 , δ): 4.15 (2H, s), 7.22 - 7.24 (2H, m), 7.31 (1H, dd, $J = 2.0$ and 8.3 Hz), 7.41 (1H, d, $J = 8.7$ Hz), 7.59 (1H, d, $J = 2.0$ Hz), 7.71 (1H, dd, $J = 1.4$ and 8.7 Hz), 8.12 (1H, brs), 11.30 (1H, brs), 12.39 (1H, brs).

<Example 19; Production of 5-(1-butanefonylcarbonyl)-3-(2,4-dichlorobenzyl)indole (48)>

According to the method of Example 3, obtained is 5-(1-butanefonylcarbonyl)-3-(2,4-dichlorobenzyl)indole (48) (0.19 g) from 5-carboxy-3-(2,4-dichlorobenzyl)indole (0.96 g), N,N'-carbonyldiimidazole (0.73 g), 1-butanefonylamine (0.62 g) and diazabicycloundecene (0.69 g).

$^1\text{H-NMR}$ (DMSO-d_6 , δ): 0.86 (3H, t, $J = 7.3$ Hz), 1.40 (2H, m), 1.67 (2H, m), 3.51 (2H, t, $J = 7.7$ Hz), 4.16 (2H, s), 7.21 (1H, d, $J = 2.2$ Hz), 7.25 (1H, d, $J = 8.3$ Hz), 7.33 (1H, dd, $J = 2.1$ and 8.3 Hz), 7.44 (1H, d, $J = 8.6$ Hz), 7.60 (1H, d, $J = 2.1$ Hz), 7.71 (1H, dd, $J = 1.7$ and 8.6 Hz), 8.28 (1H, d, $J = 1.2$ Hz), 11.37 (1H, brs), 11.77 (1H, s).

IR (KBr): 1662 cm^{-1} .

m.p.: $165.5 - 166.2^\circ\text{C}$.

Mass (FD): m/e 438 (M).

<Example 20; Production of 1-(2,4-dichlorobenzyl)-5-methoxycarbonylindole (49)>

2,4-Dichlorobenzyl chloride (8.68 g) is added to a mixture of 5-methoxycarbonylindole (6.49 g), 60 % sodium hydride (2.24 g) and N,N-dimethylformamide, while cooled in an ice-water bath. This is stirred at room temperature for 20 minutes, and water is added thereto. Then, this is extracted with ethyl acetate. After washed with water, the organic layer is dried, concentrated, purified through silica gel column chromatography (eluent: hexane/ethyl acetate = 6/1), and finally recrystallized from ethyl acetate/hexane to obtain 1-(2,4-dichlorobenzyl)-5-methoxycarbonylindole (49) (6.59 g).

¹H-NMR (CDCl₃, δ): 3.93 (3H, s), 5.39 (2H, s), 6.49 (1H, d, J = 8.4 Hz), 6.69 (1H, d, J = 3.2 Hz), 7.08 (1H, dd, J = 2.0 and 8.4 Hz), 7.17 (1H, d, J = 3.2 Hz), 7.23 (1H, d, J = 8.6 Hz), 7.45 (1H, d, J = 2.0 Hz), 7.89 (1H, dd, J = 1.4 and 8.6 Hz), 8.43 (1H, s).

<Example 21; Production of 5-carboxy-1-(2,4-dichlorobenzyl)indole (50)>

According to the method of Example 2, obtained is 5-carboxy-1-(2,4-dichlorobenzyl)indole (50) (4.52 g) from 1-(2,4-dichlorobenzyl)-5-methoxycarbonylindole (5.02 g).

¹H-NMR (DMSO-d₆, δ): 5.53 (2H, s), 6.64 (1H, d, J = 8.4 Hz), 6.68 (1H, d, J = 3.1 Hz), 7.31 (1H, dd, J = 2.1 and 8.4 Hz), 7.45 (1H, d, J = 8.7 Hz), 7.53 (1H, d, J = 3.2 Hz), 7.67 (1H, d, J = 2.1 Hz), 7.73 (1H, dd, J = 1.5 and 8.7 Hz), 8.27 (1H, d, J = 1.3 Hz), 12.48 (1H, brs).

<Example 22; Production of 5-(1-butanefonylcarbonyl)-1-(2,4-dichlorobenzyl)indole (51)>

According to the method of Example 3, obtained is 5-(1-butanefonylcarbonyl)-1-(2,4-dichlorobenzyl)indole (51) (0.44 g) from 5-carboxy-1-

(2,4-dichlorobenzyl)indole (0.96 g), N,N'-carbonyldiimidazole (0.73 g), 1-butanefulfonamide (0.62 g) and diazabicycloundecene (0.69 g).

$^1\text{H-NMR}$ (DMSO- d_6 , δ): 0.86 (3H, t, $J = 7.4$ Hz), 1.40 (2H, m), 1.68 (2H, m), 3.51 (2H, t, $J = 7.7$ Hz), 5.55 (2H, s), 6.64 (1H, d, $J = 8.4$ Hz), 6.70 (1H, d, $J = 3.2$ Hz), 7.32 (1H, dd, $J = 2.1$ and 8.4 Hz), 7.51 (1H, d, $J = 8.7$ Hz), 7.56 (1H, d, $J = 3.2$ Hz), 7.68 (1H, d, $J = 2.1$ Hz), 7.72 (1H, dd, $J = 1.6$ and 8.7 Hz), 8.31 (1H, d, $J = 1.5$ Hz), 11.83 (1H, s).

IR (KBr): 1646 cm^{-1} .

m.p.: 143.8 - 144.8°C.

<Production Example 7; methyl 3-(2-oxopropyl)-4-nitrobenzoate>

According to the method of Production Example 1, a mixed solution of methyl E-3-(2-dimethylaminovinyl)-4-nitrobenzoate (45.05 g), pyridine (23.51 g), acetyl chloride (21.20 g) and methylene chloride (200 ml) is stirred at room temperature for 61 hours, and then heated under reflux with 1,4-dioxane in water for 18 hours to obtain methyl 3-(2-oxopropyl)-4-nitrobenzoate (19.67 g).

$^1\text{H-NMR}$ (CDCl_3 , δ): 2.34 (3H, s), 3.96 (3H, s), 4.18 (2H, s), 7.95 (1H, d, $J = 1.8$ Hz), 8.10 (1H, dd, $J = 1.8$ and 8.5 Hz), 8.15 (1H, d, $J = 8.5$ Hz).

<Production Example 8; 5-methoxycarbonyl-2-methylindole>

According to the method of Production Example 5, obtained is 5-methoxycarbonyl-2-methylindole (8.25 g) from methyl 3-(2-oxopropyl)-4-nitrobenzoate (18.98 g) and sodium hydrosulfite (208.92 g).

[Physical Properties of the Product]

$^1\text{H-NMR}$ (CDCl_3 , δ): 2.46 (3H, s), 3.92 (3H, s), 6.31 (1H, s), 7.28 (1H, d, $J = 8.5$ Hz), 7.83 (1H, dd, $J = 1.5$ and 8.5 Hz), 8.08 (1H, brs), 8.27 (1H, s).

<Example 23; Production of 3-(2,4-dichlorobenzyl)-5-methoxycarbonyl-2-methylindole>

(52)>

According to the method of Example 17, obtained is 3-(2,4-dichlorobenzyl)-5-methoxycarbonyl-2-methylindole (52) (6.97 g) from 5-methoxycarbonyl-2-methylindole (7.57 g), 2,4-dichlorobenzyl iodide (13.77 g) and silver oxide (11.33 g).

$^1\text{H-NMR}$ (CDCl_3 , δ): 2.36 (3H, s), 3.89 (3H, s), 4.11 (2H, s), 6.83 (1H, d, $J = 8.3$ Hz), 7.03 (1H, dd, $J = 2.1$ and 8.3 Hz), 7.31 (1H, d, $J = 8.6$ Hz), 7.41 (1H, d, $J = 2.1$ Hz), 7.85 (1H, dd, $J = 1.5$ and 8.6 Hz), 8.09 (1H, s), 8.11 (1H, brs).

<Example 24; Production of 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (53)>

According to the method of Example 3, obtained is 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (53) (0.77 g) from 3-(2,4-dichlorobenzyl)-5-methoxycarbonyl-2-methylindole (1.74 g).

$^1\text{H-NMR}$ (DMSO-d_6 , δ): 2.34 (3H, s), 4.08 (2H, s), 6.98 (1H, d, $J = 8.4$ Hz), 7.26 (1H, dd, $J = 2.1$ and 8.4 Hz), 7.31 (1H, d, $J = 8.5$ Hz), 7.59 (1H, d, $J = 2.1$ Hz), 7.62 (1H, dd, $J = 1.5$ and 8.5 Hz), 7.87 (1H, s), 11.29 (1H, s), 12.32 (1H, s).

<Example 25; Production of 5-(1-butanefonylcarbonyl)-3-(2,4-dichlorobenzyl)-2-methylindole (54)>

According to the method of Example 3, obtained is 5-(1-butanefonylcarbonyl)-3-(2,4-dichlorobenzyl)-2-methylindole (54) (0.49 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.77 g), N,N' -carbonyldiimidazole (0.57 g), 1-butanefonyl (0.48 g) and diazabicycloundecene (0.53 g).

$^1\text{H-NMR}$ (DMSO-d_6 , δ): 0.84 (3H, t, $J = 7.4$ Hz), 1.38 (2H, m), 1.65 (2H, m), 2.30 (3H, s), 3.48 (2H, t, $J = 7.7$ Hz), 4.08 (2H, s), 6.93 (1H, d, $J = 8.4$ Hz), 7.26 (1H, dd, $J = 2.2$ and 8.4 Hz), 7.35 (1H, d, $J = 8.5$ Hz), 7.60 (1H, d, $J = 2.2$ Hz), 7.63 (1H, dd, $J = 1.7$ and 8.5 Hz), 8.04 (1H, s), 11.39 (1H, s), 11.68 (1H, s).

IR (Nujol): 1673 cm^{-1} .

m.p.: 174.9 - 175.4°C.

<Example 26; Production of 3-(2,4-dichlorobenzyl)-5-methoxycarbonyl-1-methylindole (55)>

60 % sodium hydride (0.07 g) and then methyl iodide (0.18 g) are added in that order to an N,N-dimethylformamide solution of 3-(2,4-dichlorobenzyl)-5-methoxycarbonylindole (0.34 g), and stirred at room temperature for 19 hours. After concentrated under reduced pressure, this is extracted with water and ethyl acetate. The resulting extract is washed with water, dried, concentrated, and purified through silica gel column chromatography (eluent: hexane/ethyl acetate = 9/1 to 7/1) to obtain 3-(2,4-dichlorobenzyl)-5-methoxycarbonyl-1-methylindole (55) (0.30 g).

[Physical Properties of Compound (55)]

$^1\text{H-NMR}$ (CDCl_3 , δ): 3.76 (3H, s), 3.92 (3H, s), 4.17 (2H, s), 6.80 (1H, s), 7.09 - 7.13 (2H, m), 7.30 (1H, d, $J = 8.4$ Hz), 7.41 (1H, d, $J = 1.8$ Hz), 7.93 (1H, dd, $J = 1.6$ and 8.7 Hz), 8.30 (1H, d, $J = 1.2$ Hz).

<Example 27; Production of 5-carboxy-3-(2,4-dichlorobenzyl)-1-methylindole (56)>

According to the method of Example 3, obtained is 5-carboxy-3-(2,4-dichlorobenzyl)-1-methylindole (56) (0.83 g) from 3-(2,4-dichlorobenzyl)-5-methoxycarbonyl-1-methylindole (0.89 g).

$^1\text{H-NMR}$ (DMSO-d_6 , δ): 3.76 (3H, s), 4.14 (2H, s), 7.17 (1H, s), 7.26 (1H, d, $J = 8.3$ Hz), 7.33 (1H, dd, $J = 2.1$ and 8.3 Hz), 7.46 (1H, d, $J = 8.7$ Hz), 7.60 (1H, d, $J = 2.1$ Hz), 7.75 (1H, dd, $J = 1.5$ and 8.7 Hz), 8.13 (1H, d, $J = 1.1$ Hz), 12.45 (1H, brs).

<Example 28; Production of 5-butanefonylcarbamoyl-3-(2,4-dichlorobenzyl)-1-methylindole (57)>

According to the method of Example 3, obtained is 5-(1-

butanesulfonylcarbamoyl)-3-(2,4-dichlorobenzyl)-1-methylindole (57) (0.80 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-1-methylindole (0.80 g), N,N'-carbonyldiimidazole (0.58 g), 1-butanesulfonamide (0.49 g) and diazabicycloundecene (0.56 g).

$^1\text{H-NMR}$ (DMSO- d_6 , δ): 0.86 (3H, t, $J = 7.4$ Hz), 1.40 (2H, m), 1.68 (2H, m), 3.52 (2H, t, $J = 7.7$ Hz), 3.76 (3H, s), 4.15 (2H, s), 7.16 (1H, s), 7.28 (1H, d, $J = 8.3$ Hz), 7.34 (1H, dd, $J = 2.0$ and 8.3 Hz), 7.51 (1H, d, $J = 8.8$ Hz), 7.60 (1H, d, $J = 2.0$ Hz), 7.77 (1H, dd, $J = 1.4$ and 8.8 Hz), 8.30 (1H, d, $J = 1.1$ Hz), 11.80 (1H, s).

IR (Nujol): 1680 cm^{-1} .

Foamy.

<Production Example 9; 6-methoxycarbonyl-2-methylindole>

A mixed solution of methyl 4-(2-oxopropyl)-3-nitrobenzoate (3.86 g), reduced iron (9.0 g) and acetic acid (40 ml) is stirred at 90°C for 24 hours. The solid is taken out through filtration, and the filtrate is concentrated under reduced pressure. The resulting residue is purified through silica gel column chromatography (eluent: hexane/ethyl acetate = 3/1) to obtain 6-methoxycarbonyl-2-methylindole (0.890 g).

$^1\text{H-NMR}$ (CDCl_3 , δ ppm): 2.46 (3H, s), 3.90 (3H, s), 6.25 (1H, s), 7.48 (1H, d, $J = 8.3$ Hz), 7.72 (1H, dd, $J = 1.4$ and 8.3 Hz), 7.97 (1H, s), 8.39 (1H, brs).

<Production Example 10; 5-(methoxycarbonyl)-2-methylindole>

To an ethanol (10 ml) solution of methyl 4-nitro-3-(oxopropyl)benzoate (13.0 g), added is palladium-carbon (5 %, 2.6 g) in a nitrogen atmosphere. This is purged with hydrogen, and further stirred at room temperature for 3 hours and then at 60°C for 19 hours. The solid is removed through filtration, and the filtrate is concentrated. The resulting oily product is crystallized from toluene. The crystal formed is collected and dried to obtain 5-(methoxycarbonyl)-2-methylindole (8.42 g).

$^1\text{H-NMR}$ (CDCl_3 , δ ppm): 2.45 (3H, s), 3.92 (3H, s), 6.30 (1H, d, $J = 0.7$ Hz), 7.27 (1H, d, $J = 8.3$ Hz), 7.53 (1H, dd, $J = 8.5$ and 1.6 Hz), 8.14 (1H, brs), 8.27 (1H, s).

<Production Example 11; 5-(methoxycarbonyl)-2-propylindole>

Methyl 4-nitro-3-(2-oxopentyl)benzoate (5.42 g) is dissolved in a mixed solvent of tetrahydrofuran (20 ml) and methanol (20 ml), to which is added a suspension of sodium hydrosulfite (52.2 g) in water (60 ml). After having been refluxed for 2 hours, this is subjected to liquid-liquid separation by adding ethyl acetate (300 ml) and water (200 ml) thereto. The organic layer is washed with a saline solution, dried and concentrated. The resulting residue is purified through silica gel column chromatography to obtain 5-(methoxycarbonyl)-2-propylindole (3.16 g).

$^1\text{H-NMR}$ (CDCl_3 , δ ppm): 1.02 (3H, t, $J = 7.5$ Hz), 1.77 (2H, m), 2.75 (2H, t, $J = 7.6$ Hz), 3.92 (3H, s), 6.33 (1H, dd, $J = 0.8$ and 1.7 Hz), 7.29 (1H, d, $J = 8.5$ Hz), 7.83 (1H, dd, $J = 1.6$ and 8.5 Hz), 8.07 (1H, brs), 8.28 (1H, d, $J = 1.2$ Hz).

<Production Example 12; 2-ethyl-5-(methoxycarbonyl)indole>

According to the method of Production Example 11, obtained is 2-ethyl-5-(methoxycarbonyl)indole (1.88 g) from methyl 4-nitro-3-(2-oxobutyl)benzoate (3.77 g) and sodium hydrosulfite (39.17 g).

$^1\text{H-NMR}$ (CDCl_3 , δ ppm): 1.36 (3H, t, $J = 7.6$ Hz), 2.81 (2H, q, $J = 7.6$ Hz), 3.92 (3H, s), 6.33 (1H, d, $J = 0.8$ Hz), 7.29 (1H, d, $J = 8.5$ Hz), 7.83 (1H, dd, $J = 1.6$ and 8.5 Hz), 8.08 (1H, brs), 8.29 (1H, s).

<Production Example 13; 5-(methoxycarbonyl)indole>

A mixture of 5-carboxyindole (16.0 g), sodium hydrogencarbonate (26.68 g), methyl iodide (61.11 g) and N,N -dimethylformamide (30 ml) is stirred at room temperature for 4 days. Water and ethyl acetate are added thereto for liquid-liquid

separation. The organic layer is washed with aqueous sodium hydrogencarbonate, dried and concentrated to obtain 5-(methoxycarbonyl)indole (13.74 g).

$^1\text{H-NMR}$ (CDCl_3 , δ ppm): 3.93 (3H, s), 6.65 (1H, s), 7.27 (1H, m), 7.40 (1H, d, $J = 8.6$ Hz), 7.91 (1H, dd, $J = 8.6$ and 1.6 Hz), 8.42 (1H, s), 8.46 (1H, brs).

<Production Example 14: 6-(methoxycarbonyl)-3-methylindole>

Methyl 4-ethyl-3-nitrobenzoate (5.0 g), which had been prepared through nitration followed by methyl-esterification of 4-ethylbenzoic acid, is dissolved in N,N-dimethylformamide (50 ml), to which is added N,N-dimethylformamide dimethyl acetal (8.45 g), and stirred under heat at 130°C for 3 hours. The reaction mixture is concentrated under reduced pressure, and the resulting red oily residue is dissolved in methanol (50 ml), to which is added palladium-carbon (5 %, 0.400 g), and stirred in a hydrogen atmosphere at room temperature for 2.5 hours. Then, this is further stirred at 50°C for 2 hours and thereafter at room temperature for 3 days. The catalyst is removed through filtration, and the filtrate is concentrated under reduced pressure. The resulting residue is recrystallized from t-butyl methyl ether to obtain a pale yellow crystal of the intended product (2.4 g).

$^1\text{H-NMR}$ (CDCl_3 , δ ppm): 8.17 (1H, br), 8.10 (1H, d, $J = 1.3$ Hz), 7.81 (1H, dd, $J = 1.3$ and 8.3 Hz), 7.59 (1H, d, $J = 8.3$ Hz), 7.13 (1H, q, $J = 0.9$ Hz), 3.93 (3H, s), 2.34 (3H, d, $J = 0.9$ Hz).

<Example 29; Production of 3-(2,4-dichlorobenzyl)-6-(methoxycarbonyl)-2-methylindole (58)>

6-(Methoxycarbonyl)-2-methylindole (3.03 g), 2,4-dichlorobenzyl chloride (4.69 g) and silver(I) oxide (5.56 g) are suspended in 1,4-dioxane (50 ml), and stirred under heat at 90°C for 19.5 hours. The solid is separated through filtration, and the filtrate is concentrated. The resulting residue is purified through silica gel column

chromatography to obtain 3-(2,4-dichlorobenzyl)-6-(methoxycarbonyl)-2-methylindole (58) (1.15 g).

$^1\text{H-NMR}$ (CDCl_3 , δ ppm): 2.40 (3H, s), 3.92 (3H, s), 4.10 (2H, s), 6.84 (1H, d, J = 8.3 Hz), 7.04 (1H, dd, J = 2.1 and 8.4 Hz), 7.31 (1H, d, J = 8.3 Hz), 7.40 (1H, d, J = 2.1 Hz), 7.73 (1H, dd, J = 1.4 and 8.3 Hz), 8.06 (1H, d, J = 1.1 Hz), 8.14 (1H, brs).

<Example 30; Production of 3-(biphenyl-4-ylmethyl)-5-(methoxycarbonyl)-2-methylindole (59)>

According to the method of Example 29, obtained is 3-(biphenyl-4-ylmethyl)-5-(methoxycarbonyl)-2-methylindole (59) (0.386 g) from 5-(methoxycarbonyl)-2-methylindole (0.946 g), 4-(bromomethyl)biphenyl (1.73 g) and silver(I) oxide (1.62 g).

$^1\text{H-NMR}$ (CDCl_3 , δ ppm): 2.42 (3H, s), 3.89 (3H, s), 4.14 (2H, s), 7.26 - 7.32 (4H, m), 7.40 (2H, t, J = 7.5 Hz), 7.47 (2H, d, J = 8.2 Hz), 7.54 (2H, m), 7.84 (1H, dd, J = 1.4 and 8.6 Hz), 8.01 (1H, brs), 8.23 (1H, s).

<Example 31; Production of 3-(2-chlorobenzyl)-5-(methoxycarbonyl)-2-methylindole (60)>

According to the method of Example 29, obtained is 3-(2-chlorobenzyl)-5-(methoxycarbonyl)-2-methylindole (60) (0.18 g) from 5-(methoxycarbonyl)-2-methylindole (0.946 g), 2-chlorobenzyl bromide (1.43 g) and silver(I) oxide (1.62 g).

$^1\text{H-NMR}$ (CDCl_3 , δ ppm): 2.36 (3H, s), 3.88 (3H, s), 4.17 (2H, s), 6.93 (1H, d, J = 7.8 Hz), 7.06 (1H, dt, J = 1.3 and 7.5 Hz), 7.12 (1H, dt, J = 1.5 and 7.6 Hz), 7.30 (1H, d, J = 8.4 Hz), 7.39 (1H, dd, J = 1.3 and 7.9 Hz), 7.84 (1H, dd, J = 1.5 and 8.4 Hz), 8.06 (1H, brs), 8.13 (1H, s).

<Example 32; Production of 3-(2,4-dichlorobenzoyl)-5-(methoxycarbonyl)-2-methylindole (61)>

Aluminium chloride (0.86 g) is suspended in methylene chloride (30 ml), to which are added 2,4-dichlorobenzoyl chloride (0.67 g) and then 5-(methoxycarbonyl)-2-methylindole (0.50 g), and stirred at 50°C for 3 hours and then at room temperature for 41 hours. The reaction mixture is poured into water with ice, and extracted twice with ethyl acetate. The organic layer is washed twice with a saturated aqueous sodium hydrogencarbonate solution and then with a saturated saline solution, and thereafter dried with anhydrous sodium sulfate. The drying agent is removed through filtration, and the filtrate is concentrated under reduced pressure. The resulting crystal is washed with t-butyl methyl ether, and dried to obtain a crystal of 3-(2,4-dichlorobenzoyl)-5-(methoxycarbonyl)-2-methylindole (61) (0.60 g).

¹H-NMR (DMSO-d₆, δ ppm): 12.45 (1H, br), 8.20 (1H, s), 7.81 (1H, d, J = 1.9 Hz), 7.79 (1H, dd, J = 1.7 and 8.4 Hz), 7.60 (1H, dd, J = 1.9 and 8.3 Hz), 7.48 (1H, d, J = 8.1 Hz), 7.47 (1H, d, J = 8.3 Hz), 3.81 (3H, s), 2.23 (3H, s).

<Example 33; Production of 3-(4-benzyloxybenzyl)-5-(methoxycarbonyl)-2-methylindole (62)>

A methylene chloride (15 ml) solution of 5-(methoxycarbonyl)-2-methylindole (0.567 g) and 4-benzyloxybenzaldehyde (0.700 g) is dropwise added to a methylene chloride (15 ml) solution of trifluoroacetic acid (0.513 g) and triethylsilane (1.047 g) in an ice-water bath, over a period of 15 minutes. This is stirred for 30 minutes in the ice-water bath, and an aqueous solution of 2 M sodium hydroxide is added thereto to stop the reaction. The organic layer is washed with a saturated saline solution, dried and concentrated. The resulting residue is purified through silica gel column chromatography (eluent: ethyl acetate/hexane = 3/7) to obtain 3-(4-benzyloxybenzyl)-5-(methoxycarbonyl)-2-methylindole (62) (0.510 g).

¹H-NMR (CDCl₃, δ ppm): 2.34 (3H, s), 3.88 (3H, s), 4.01 (2H, s), 4.98 (2H, s),

6.84 (2H, d, J = 8.6 Hz), 7.09 (2H, d, J = 8.6 Hz), 7.23 (1H, d, J = 8.3 Hz), 7.30 (1H, d, J = 7.2 Hz), 7.35 (2H, m), 7.39 (2H, m), 7.81 (1H, dd, J = 1.5 and 8.4 Hz), 8.12 (1H, brs), 8.19 (1H, s).

<Example 34: Production of 3-(2,4-dichlorobenzyl)-5-(methoxycarbonyl)-2-propylindole (63)>

According to the method of Example 33, obtained is 3-(2,4-dichlorobenzyl)-5-(methoxycarbonyl)-2-propylindole (63) (0.617 g) from 5-(methoxycarbonyl)-2-propylindole (0.652 g), 2,4-dichlorobenzaldehyde (0.578 g), trifluoroacetic acid (0.513 g) and triethylsilane (1.047 g).

¹H-NMR (CDCl₃, δ ppm): 0.93 (3H, t, J = 7.4 Hz), 1.64 (2H, m), 2.68 (2H, t, J = 7.7 Hz), 3.88 (3H, s), 4.12 (2H, s), 6.79 (1H, d, J = 8.3 Hz), 7.01 (1H, dd, J = 2.0 and 8.4 Hz), 7.32 (1H, d, J = 8.5 Hz), 7.41 (1H, d, J = 2.0 Hz), 7.86 (1H, d, J = 8.3 Hz), 8.07 (1H, s), 8.10 (1H, brs).

<Example 35: Production of 3-(2,4-dichlorobenzyl)-2-ethyl-5-(methoxycarbonyl)indole (64)>

According to the method of Example 33, obtained is 3-(2,4-dichlorobenzyl)-2-ethyl-5-(methoxycarbonyl)indole (64) (0.420 g) from 2-ethyl-5-(methoxycarbonyl)indole (0.610 g), 2,4-dichlorobenzaldehyde (0.578 g), trifluoroacetic acid (0.513 g) and triethylsilane (1.047 g).

¹H-NMR (CDCl₃, δ ppm): 1.24 (3H, t, J = 7.6 Hz), 2.73 (2H, q, J = 7.6 Hz), 3.89 (3H, s), 4.17 (2H, s), 6.80 (1H, d, J = 8.4 Hz), 7.01 (1H, dd, J = 2.1 and 8.4 Hz), 7.33 (1H, d, J = 8.5 Hz), 7.41 (1H, d, J = 2.1 Hz), 7.86 (1H, dd, J = 1.5 and 8.5 Hz), 8.08 (1H, s), 8.14 (1H, brs).

<Example 36: Production of 3-(2,4-dichlorobenzyl)-5-(methoxycarbonyl)indole (65)>

From (5-methoxycarbonyl)indole (17.52 g), 2,4-dichlorobenzyl iodide (34.49 g)

and silver oxide (28.09 g), obtained is 3-(2,4-dichlorobenzyl)-5-(methoxycarbonyl)indole (65) (5.49 g).

$^1\text{H-NMR}$ (CDCl_3 , δ ppm): 3.92 (3H, s), 4.20 (2H, s), 6.98 (1H, d, $J = 2.3$ Hz), 7.09 - 7.13 (2H, m), 7.38 (1H, d, $J = 9.1$ Hz), 7.42 (1H, d, $J = 1.9$ Hz), 7.92 (1H, dd, $J = 1.5$ and 8.5 Hz), 8.22 (1H, brs), 8.31 (1H, d, $J = 1.2$ Hz).

<Example 37; Production of 3-(1-bromonaphthalen-2-ylmethyl)-5-(methoxycarbonyl)-2-methylindole (66)>

According to the method of Example 33, obtained is 3-(1-bromonaphthalen-2-ylmethyl)-5-(methoxycarbonyl)-2-methylindole (66) (0.339 g) from 5-(methoxycarbonyl)-2-methylindole (0.73 g), 1-bromo-2-naphthaldehyde (1.00 g), trifluoroacetic acid (0.662 g) and triethylsilane (1.35 g).

$^1\text{H-NMR}$ (CDCl_3 , δ ppm): 2.36 (3H, s), 3.86 (3H, s), 4.43 (2H, s), 7.08 (1H, d, $J = 8.5$ Hz), 7.31 (1H, d, $J = 8.4$ Hz), 7.47 (1H, t, $J = 8.1$ Hz), 7.58 - 7.61 (2H, m), 7.75 (1H, d, $J = 8.5$ Hz), 7.85 (1H, dd, $J = 8.5$ and 1.6 Hz), 8.05 (1H, brs), 8.18 (1H, s), 8.38 (1H, d, $J = 8.6$ Hz).

<Example 38; Production of 3-((3-chloropyridin-4-yl)methyl)-5-(methoxycarbonyl)-2-methylindole (67)>

According to the method of Example 33, obtained is 3-((3-chloropyridin-4-yl)methyl)-5-(methoxycarbonyl)-2-methylindole (67) (0.355 g) from 5-(methoxycarbonyl)-2-methylindole (0.486 g), 3-chloropyridine-4-carboxyaldehyde (0.40 g), triethylsilane (0.896 g) and trifluoroacetic acid (0.439 g).

$^1\text{H-NMR}$ (CDCl_3 , δ ppm): 2.38 (3H, s), 3.89 (3H, s), 4.16 (2H, s), 6.82 (1H, d, $J = 5.0$ Hz), 7.33 (1H, d, $J = 8.5$ Hz), 7.87 (1H, d, $J = 8.7$ Hz), 8.08 (1H, s), 8.20 (1H, brs), 8.25 (1H, d, $J = 5.0$ Hz), 8.57 (1H, s).

<Example 39; Production of 5-(methoxycarbonyl)-2-methyl-3-(4-(2-

phenylethenyl)benzyl)indole (68)>

According to the method of Example 33, obtained is 5-(methoxycarbonyl)-2-methyl-3-(4-(2-phenylethenyl)benzyl)indole (68) (0.16 g) from 5-(methoxycarbonyl)-2-methylindole (0.567 g), 4-stilbenecarboxyaldehyde (0.687 g), trifluoroacetic acid (0.513 g) and triethylsilane (1.047 g).

$^1\text{H-NMR}$ (CDCl_3 , δ ppm): 2.40 (3H, s), 3.89 (3H, s), 4.10 (2H, s), 7.05 (2H, d, J = 4.5 Hz), 7.19 (2H, d, J = 8.2 Hz), 7.23 (1H, tt, J = 1.0 and 7.4 Hz), 7.29 (1H, d, J = 8.9 Hz), 7.34 (2H, t, J = 7.7 Hz), 7.40 (2H, d, J = 8.2 Hz), 7.48 (2H, m), 7.84 (1H, dd, J = 1.6 and 8.4 Hz), 7.99 (1H, brs), 8.20 (1H, s).

<Example 40; Production of 3-((4-chloroisoquinolin-3-yl)methyl)-5-(methoxycarbonyl)-2-methylindole (69)>

To a mixture of L-tartaric acid (0.600 g), sodium hydroxide (0.160 g), 1,4-dioxane (4 ml) and water (4 ml), added are sodium iodide (0.15 g) and 5-(methoxycarbonyl)-2-methylindole (0.378 g). Then, a 1,4-dioxane (2 ml) solution of 4-chloro-3-(chloromethyl)isoquinoline (0.50 g) is added thereto, and stirred at 100°C for 56 hours. The reaction mixture is concentrated, and water is added to the residue, which is then extracted with ethyl acetate. The organic layer is dried and concentrated, and the resulting residue is purified through silica gel column chromatography to obtain 3-((4-chloroisoquinolin-3-yl)methyl)-5-(methoxycarbonyl)-2-methylindole (69) (0.14 g).

$^1\text{H-NMR}$ (CDCl_3 , δ ppm): 2.50 (3H, s), 3.89 (3H, s), 4.58 (2H, s), 7.22 (1H, d, J = 8.4 Hz), 7.59 (1H, m), 7.75 - 7.79 (2H, m), 7.93 (1H, d, J = 8.1 Hz), 7.98 (1H, brs), 8.22 (1H, d, J = 8.4 Hz), 8.45 (1H, s), 9.11 (1H, s).

<Example 41; Production of 3-((4-bromoisoquinolin-3-yl)methyl)-5-(methoxycarbonyl)-2-methylindole (70)>

According to the method of Example 40, obtained is 3-((4-bromoisquinolin-3-yl)methyl)-5-(methoxycarbonyl)-2-methylindole (70) (0.178 g) from 5-(methoxycarbonyl)-2-methylindole (0.687 g), 4-bromo-3-(bromomethyl)isoquinoline (1.31 g), L-tartaric acid (1.088 g), sodium hydroxide (0.290 g) and sodium iodide (0.217 g).

$^1\text{H-NMR}$ (CDCl_3 , δ ppm): 2.32 (3H, s), 3.86 (3H, s), 4.60 (2H, s), 7.07 (1H, d, $J = 8.4$ Hz), 7.54 (1H, t, $J = 7.6$ Hz), 7.72 (2H, m), 7.82 (1H, d, $J = 8.2$ Hz), 8.19 (1H, d, $J = 8.5$ Hz), 8.37 (1H, s), 8.51 (1H, brs), 9.03 (1H, s).

<Example 42; Production of 2-(2,4-dichlorobenzoyl)-6-(methoxycarbonyl)-3-methylindole (71)>

Aluminium chloride (1.72 g) is suspended in methylene chloride (50 ml), to which are added 2,4-dichlorobenzoyl chloride (1.35 g) and then 6-(methoxycarbonyl)-3-methylindole (1.00 g), and stirred at room temperature for 16 hours and then at 40°C for 2.5 hours. The reaction mixture is poured into water with ice, and then extracted with ethyl acetate (150 ml). The organic layer is washed three times with a saturated aqueous sodium hydrogencarbonate solution (50 ml) and then with a saturated saline solution, and thereafter dried with anhydrous sodium sulfate. The drying agent is removed through filtration, and the filtrate is concentrated under reduced pressure. The resulting crystal is washed with methyl t-butyl ether and dried to obtain 2-(2,4-dichlorobenzoyl)-6-(methoxycarbonyl)-3-methylindole (71) (1.48 g).

$^1\text{H-NMR}$ (CDCl_3 , δ ppm): 2.07 (3H, s), 3.96 (3H, s), 7.36 (1H, d, $J = 8.2$ Hz), 7.42 (1H, dd, $J = 1.9$ and 8.3 Hz), 7.54 (1H, d, $J = 1.8$ Hz), 7.69 (1H, d, $J = 8.7$ Hz), 7.81 (1H, dd, $J = 1.4$ and 8.6 Hz), 8.15 (1H, s), 9.12 (1H, brs).

<Example 43; Production of 1-(2,4-dichlorobenzyl)-6-methoxycarbonyl)-3-methylindole (72)>

6-(Methoxycarbonyl)-3-methylindole (0.57 g) is dissolved in N,N-dimethylformamide (10 ml), to which is added sodium hydride (60 % oily, 0.145 g) with cooling with ice. Then, 2,4-dichlorobenzyl chloride (0.707 g) is added thereto, and stirred at room temperature for 1.5 hours. Water is added to the reaction mixture, which is then extracted with ethyl acetate. The organic layer is washed with a saturated saline solution, and dried with anhydrous magnesium sulfate. The drying agent is removed through filtration, and the filtrate is concentrated under reduced pressure. The resulting residue is purified through silica gel column chromatography (eluent: hexane/ethyl acetate = 9/1) to obtain 1-(2,4-dichlorobenzyl)-6-(methoxycarbonyl)-3-methylindole (72) (0.83 g).

$^1\text{H-NMR}$ (CDCl_3 , δ ppm): 2.35 (3H, d, $J = 0.7$ Hz), 3.91 (3H, s), 5.36 (2H, s), 6.48 (1H, d, $J = 8.3$ Hz), 7.01 (1H, d, $J = 0.8$ Hz), 7.05 (1H, dd, $J = 2.1$ and 8.3 Hz), 7.43 (1H, d, $J = 2.0$ Hz), 7.61 (1H, d, $J = 8.3$ Hz), 7.82 (1H, dd, $J = 1.3$ and 8.3 Hz), 7.96 (1H, s).

<Example 44; Production of 6-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (73)>

A mixture of 3-(2,4-dichlorobenzyl)-6-(methoxycarbonyl)-2-methylindole (0.330 g), ethanol (15 ml) and 10 % aqueous sodium hydroxide (10 ml) is heated under reflux for 1.5 hours. This is processed with hydrochloric acid to have a pH of 3, and the crystal formed is taken out through filtration. This is dried to obtain 6-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (73) (0.305 g).

[Physical Properties of Compound (73)]

$^1\text{H-NMR}$ (DMSO-d_6 , ppm): 2.35 (3H, s), 4.06 (2H, s), 7.01 (1H, d, $J = 8.4$ Hz), 7.26 (2H, m), 7.49 (1H, d, $J = 8.3$ Hz), 7.57 (1H, d, $J = 2.0$ Hz), 7.89 (1H, s), 11.38 (1H, s).

<Example 45; Production of 3-(biphenyl-4-ylmethyl)-5-carboxy-2-methylindole (74)>

According to the method of Example 44, obtained is 3-(biphenyl-4-ylmethyl)-5-carboxy-2-methylindole (74) (0.359 g) from 3-(biphenyl-4-ylmethyl)-5-(methoxycarbonyl)-2-methylindole (0.381 g).

¹H-NMR (DMSO-d₆, δ ppm): 2.37 (3H, s), 4.06 (2H, s), 7.26 (2H, d, J = 8.1 Hz), 7.30 (2H, m), 7.41 (2H, t, J = 7.6 Hz), 7.52 (2H, d, J = 8.1 Hz), 7.58 (2H, d, J = 7.7 Hz), 7.61 (1H, dd, J = 1.5 and 8.5 Hz), 8.01 (1H, s), 11.24 (1H, s), 12.29 (1H, brs).

<Example 46; Production of 5-carboxy-3-(2-chlorobenzyl)-2-methylindole (75)>

According to the method of Example 44, obtained is 5-carboxy-3-(2-chlorobenzyl)-2-methylindole (75) (0.164 g) from 3-(2-chlorobenzyl)-5-(methoxycarbonyl)-2-methylindole (0.179 g).

¹H-NMR (DMSO-d₆, δ ppm): 2.34 (3H, s), 4.10 (2H, s), 7.00 (1H, dd, J = 2.0 and 7.6 Hz), 7.18 (2H, m), 7.30 (1H, d, J = 8.4 Hz), 7.43 (1H, dd, J = 1.6 and 7.5 Hz), 7.61 (1H, dd, J = 1.5 and 8.4 Hz), 7.89 (1H, s), 11.25 (1H, s), 12.22 (1H, brs).

<Example 47; Production of 5-carboxy-3-(2,4-dichlorobenzoyl)-2-methylindole (76)>

3-(2,4-Dichlorobenzoyl)-5-(methoxycarbonyl)-2-methylindole (0.60 g) is suspended in ethanol (10 ml), to which is added an aqueous solution of 1.5 M sodium hydroxide (5 ml), and heated under reflux for 3 hours. The reaction mixture is concentrated under reduced pressure, and the residue is made acidic with water and 3 M HCl added thereto. The crystal formed is taken out through filtration, then suspended in ethanol-toluene, and concentrated under reduced pressure for azeotropic dehydration. Next, this is dried on calcium chloride in a vacuum desiccator to obtain a crystal of 5-carboxy-3-(2,4-dichlorobenzoyl)-2-methylindole (76) (0.56 g).

¹H-NMR (DMSO-d₆, δ ppm): 2.22 (3H, s), 7.45 (1H, d, J = 8.4 Hz), 7.48 (1H,

d, J = 8.2 Hz), 7.58 (1H, dd, J = 1.9 and 8.0 Hz), 7.78 (1H, dd, J = 1.5 and 8.4 Hz), 7.80 (1H, d, J = 1.9 Hz), 8.24 (1H, brs), 12.43 (1H, brs).

<Example 48; Production of 3-(4-benzyloxybenzyl)-5-carboxy-2-methylindole (77)>

From 3-(4-benzyloxybenzyl)-5-(methoxycarbonyl)-2-methylindole (0.500 g), obtained is 3-(4-benzyloxybenzyl)-5-carboxy-2-methylindole (77) (0.465 g).

[Physical Properties of Compound (77)]

¹H-NMR (DMSO-d₆, ppm): 2.37 (3H, s), 3.94 (2H, s), 5.01 (2H, s), 6.87 (2H, d, J = 8.6 Hz), 7.08 (2H, d, J = 8.6 Hz), 7.27 (1H, d, J = 8.5 Hz), 7.30 (1H, d, J = 7.3 Hz), 7.34 - 7.40 (4H, m), 7.60 (1H, dd, J = 1.5 and 8.5 Hz), 7.95 (1H, s), 11.15 (1H, s), 12.26 (1H, s).

<Example 49; Production of 5-carboxy-3-(2,4-dichlorobenzyl)-2-propylindole (78)>

According to the method of Example 44, obtained is 5-carboxy-3-(2,4-dichlorobenzyl)-2-propylindole (78) (0.188 g) from 3-(2,4-dichlorobenzyl)-5-(methoxycarbonyl)-2-propylindole (0.610 g).

¹H-NMR (DMSO-d₆, δ ppm): 0.85 (3H, t, J = 7.3 Hz), 1.60 (2H, m), 2.67 (2H, t, J = 7.4 Hz), 4.09 (2H, s), 6.91 (1H, d, J = 8.4 Hz), 7.25 (1H, dd, J = 2.3 and 8.4 Hz), 7.32 (1H, d, J = 8.4 Hz), 7.61 (1H, d, J = 2.1 Hz), 7.63 (1H, dd, J = 1.5 and 8.4 Hz), 7.86 (1H, s), 11.28 (1H, brs), 12.30 (1H, brs).

<Example 50; Production of 5-carboxy-3-(2,4-dichlorobenzyl)-2-ethylindole (79)>

According to the method of Example 44, obtained is 5-carboxy-3-(2,4-dichlorobenzyl)-2-ethylindole (79) (0.373 g) from 3-(2,4-dichlorobenzyl)-2-ethyl-5-(methoxycarbonyl)indole (0.410 g).

¹H-NMR (DMSO-d₆, δ ppm): 1.17 (3H, t, J = 7.5 Hz), 2.71 (2H, q, J = 7.6 Hz), 4.09 (2H, s), 6.92 (1H, d, J = 8.4 Hz), 7.26 (1H, dd, J = 2.2 and 8.3 Hz), 7.33 (1H, d, J = 8.5 Hz), 7.60 (1H, d, J = 2.1 Hz), 7.63 (1H, dd, J = 1.5 and 8.4 Hz),

7.87 (1H, d, J = 1.0 Hz), 11.33 (1H, brs), 12.29 (1H, brs).

<Example 51; Production of 5-carboxy-3-(2,4-dichlorobenzyl)indole (80)>

According to the method of Example 44, obtained is 5-carboxy-3-(2,4-dichlorobenzyl)indole (80) (4.64 g) from 3-(2,4-dichlorobenzyl)-5-(methoxycarbonyl)indole (5.02 g).

¹H-NMR (DMSO-d₆, δ ppm): 4.15 (2H, s), 7.22 - 7.24 (2H, m), 7.31 (1H, dd, J = 2.0 and 8.3 Hz), 7.41 (1H, d, J = 8.7 Hz), 7.59 (1H, d, J = 2.0 Hz), 7.71 (1H, dd, J = 1.4 and 8.7 Hz), 8.12 (1H, brs), 11.30 (1H, brs), 12.39 (1H, brs).

<Example 52; Production of 3-(1-bromonaphthalen-2-ylmethyl)-5-carboxy-2-methylindole (81)>

According to the method of Example 44, obtained is 3-(1-bromonaphthalen-2-ylmethyl)-5-carboxy-2-methylindole (81) (0.300 g) from 3-(1-bromonaphthalen-2-ylmethyl)-5-(methoxycarbonyl)-2-methylindole (0.325 g).

¹H-NMR (DMSO-d₆, δ ppm): 2.35 (3H, s), 4.34 (2H, s), 7.13 (1H, d, J = 8.5 Hz), 7.15 (1H, d, J = 8.5 Hz), 7.54 (1H, t, J = 7.1 Hz), 7.63 - 7.69 (2H, m), 7.75 (1H, d, J = 8.5 Hz), 7.88 (2H, m), 8.27 (1H, d, J = 7.7 Hz), 10.99 (1H, s).

<Example 53; Production of 5-carboxy-3-((3-chloropyridin-4-yl)methyl)-2-methylindole (82)>

According to the method of Example 44, obtained is 5-carboxy-3-((3-chloropyridin-4-yl)methyl)-2-methylindole (82) (0.380 g) from 3-((3-chloropyridin-4-yl)methyl)-5-(methoxycarbonyl)-2-methylindole (0.431 g).

¹H-NMR (DMSO-d₆, δ ppm): 2.36 (3H, s), 4.15 (2H, s), 6.99 (1H, d, J = 5.0 Hz), 7.32 (1H, d, J = 8.4 Hz), 7.64 (1H, d, J = 8.5 Hz), 7.90 (1H, s), 8.33 (1H, d, J = 4.9 Hz), 8.59 (1H, s), 11.34 (1H, s), 12.31 (1H, brs).

<Example 54; Production of 5-carboxy-2-methyl-3-(4-(2-phenylethenyl)benzyl)indole

(83)>

According to the method of Example 44, obtained is 5-carboxy-2-methyl-3-(4-(2-phenylethenyl)benzyl)indole (83) (0.095 g) from 5-(methoxycarbonyl)-2-methyl-3-(4-(2-phenylethenyl)benzyl)indole (0.16 g).

¹H-NMR (DMSO-d₆, δ ppm): 2.39 (3H, s), 4.03 (2H, s), 7.16 (2H, d, J = 7.5 Hz), 7.19 (2H, d, J = 8.1 Hz), 7.23 (1H, t, J = 7.4 Hz), 7.29 (1H, d, J = 8.4 Hz), 7.34 (2H, t, J = 7.6 Hz), 7.47 (2H, d, J = 8.0 Hz), 7.55 (2H, d, J = 7.9 Hz), 7.61 (1H, d, J = 8.4 Hz), 7.98 (1H, s), 11.22 (1H, s), 12.27 (1H, s).

<Example 55; Production of 5-carboxy-3-((4-chloroisoquinolin-3-yl)methyl)-2-methylindole (84)>

From 3-((4-chloroisoquinolin-3-yl)methyl)-5-(methoxycarbonyl)-2-methylindole (0.140 g), obtained is 5-carboxy-3-((4-chloroisoquinolin-3-yl)methyl)-2-methylindole (84) (0.112 g). This is directly used in the next reaction.

<Example 56; Production of 3-((4-bromoisoquinolin-3-yl)methyl)-5-carboxy-2-methylindole (85)>

According to the method of Example 44, obtained is 3-((4-bromoisoquinolin-3-yl)methyl)-5-carboxy-2-methylindole (85) (0.123 g) from 3-((4-bromoisoquinolin-3-yl)methyl)-5-(methoxycarbonyl)-2-methylindole (0.178 g).

¹H-NMR (DMSO-d₆, δ ppm): 2.38 (3H, s), 4.49 (2H, s), 7.04 (1H, d, J = 8.2 Hz), 7.59 (1H, d, J = 8.3 Hz), 7.70 (1H, t, J = 7.5 Hz), 7.89 (1H, t, J = 7.6 Hz), 8.07 (1H, s), 8.11 (1H, d, J = 8.3 Hz), 8.14 (1H, d, J = 8.7 Hz), 9.19 (1H, s), 10.74 (1H, s).

<Example 57; Production of 6-carboxy-2-(2,4-dichlorobenzyl)-3-methylindole (86)>

2-(2,4-Dichlorobenzoyl)-6-(methoxycarbonyl)-3-methylindole (1.00 g) is suspended in ethylene glycol (10 ml), to which is added hydrazine monohydrate (0.83

g), and stirred under heat at 160°C for 3 hours. After this is once cooled, potassium hydroxide (1.1 g) is added thereto, and again stirred under heat at 155°C for 3 hours. After cooled, the reaction mixture is made acidic with water and 1 M HCl added thereto. Then, this is extracted with ethyl acetate. The organic layer is washed with a saturated saline solution, and dried with anhydrous sodium sulfate. Then, the drying agent is removed through filtration, and the filtrate is concentrated under reduced pressure. The resulting crystalline residue is washed with a mixed solution of ethyl acetate and hexane, and dried to obtain 6-carboxy-2-(2,4-dichlorobenzyl)-3-methylindole (86) (0.63 g).

¹H-NMR (DMSO-d₆, δ ppm): 2.18 (3H, s), 4.17 (2H, s), 7.12 (1H, d, J = 8.3 Hz), 7.35 (1H, dd, J = 2.1 and 8.3 Hz), 7.47 (1H, d, J = 8.3 Hz), 7.58 (1H, dd, J = 1.0 and 8.2 Hz), 7.62 (1H, d, J = 2.1 Hz), 7.89 (1H, s), 11.02 (1H, brs).

<Example 58; Production of 6-carboxy-1-(2,4-dichlorobenzyl)-3-methylindole (87)>

According to the method of Example 44, obtained is 6-carboxy-1-(2,4-dichlorobenzyl)-3-methylindole (87) (0.46 g) from 1-(2,4-dichlorobenzyl)-6-(methoxycarbonyl)-3-methylindole (0.50 g).

¹H-NMR (DMSO-d₆, δ ppm): 12.55 (1H, brs), 7.95 (1H, s), 7.68 (1H, d, J = 2.1 Hz), 7.65 (1H, dd, J = 1.2 and 8.3 Hz), 7.59 (1H, d, J = 8.3 Hz), 7.40 (1H, s), 7.32 (1H, dd, J = 2.0 and 8.3 Hz), 6.61 (1H, d, J = 8.3 Hz), 5.51 (2H, s), 2.27 (3H, s).

<Example 59; Production of 3-(2,4-dichlorobenzyl)-5-(1-pentanesulfonylcarbamoyl)indole (88)>

N,N'-Carbonyldiimidazole (0.282 g) is added to an N,N-dimethylformamide (4.3 ml) solution of 5-carboxy-3-(2,4-dichlorobenzyl)indole (0.429 g), and stirred at room temperature for 1 hour. Diazabicycloundecene (0.306 g) and 1-pentanesulfonamide

(0.304 g) are added thereto, and stirred at 100°C for 40 hours. The reaction mixture is cooled, and then made acidic with water and 1M HCl added thereto. Then, the gummy residue formed is collected. This is dissolved in ethyl acetate, washed with water, and dried. The solvent is evaporated away, and the resulting residue is crystallized from ether. The crystal formed is collected and dried to obtain 3-(2,4-dichlorobenzyl)-5-(1-pentanesulfonylcarbamoyl)indole (88) (0.258 g).

¹H-NMR (DMSO-d₆, δ ppm): 0.83 (3H, t, J = 6.0 Hz), 1.22 - 1.43 (4H, m), 1.70 (2H, m), 3.52 (2H, t, J = 6.0 Hz), 4.18 (2H, s), 7.21 - 7.28 (2H, m), 7.35 (1H, d, J = 8.0 Hz), 7.46 (1H, d, J = 8.0 Hz), 7.62 (1H, s), 7.72 (1H, d, J = 8.0 Hz), 8.28 (1H, s), 11.38 (1H, s), 11.76 (1H, s).

m.p.: 157 - 158°C.

<Example 60; Production of 3-(2,4-dichlorobenzyl)-2-methyl-5-(1-pentanesulfonylcarbamoyl)indole (89)>

According to the method of Example 59, obtained is 3-(2,4-dichlorobenzyl)-2-methyl-5-(1-pentanesulfonylcarbamoyl)indole (89) (0.907 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (1.00 g), N,N'-carbonyldiimidazole (0.631 g), 1-pentanesulfonamide (0.588 g) and diazabicycloundecene (0.592 g).

¹H-NMR (DMSO-d₆, δ ppm): 0.80 (3H, t, J = 7.3 Hz), 1.25 (2H, m), 1.34 (2H, m), 1.67 (2H, m), 2.30 (3H, s), 3.47 (2H, t, J = 7.7 Hz), 4.08 (2H, s), 6.93 (1H, d, J = 8.4 Hz), 7.25 (1H, dd, J = 8.4 and 2.1 Hz), 7.35 (1H, d, J = 8.5 Hz), 7.60 (1H, d, J = 2.1 Hz), 7.63 (1H, dd, J = 8.5 and 1.6 Hz), 8.03 (1H, s), 11.38 (1H, s), 11.67 (1H, s).

IR (Nujol): 1682 cm⁻¹.

m.p.: 177 - 178.5°C.

<Example 61; Production of 6-(1-butanesulfonylcarbamoyl)-3-(2,4-dichlorobenzyl)-2-

methylindole (90)>

According to the method of Example 59, obtained is 6-(1-butanefonylcarbonyl)-3-(2,4-dichlorobenzyl)-2-methylindole (90) (0.133 g) from 6-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.300 g), N,N'-carbonyldiimidazole (0.292 g), diazabicycloundecene (0.274 g) and 1-butanefonyl (0.247 g).

¹H-NMR (DMSO-d₆, δ ppm): 0.86 (3H, t, J = 7.4 Hz), 1.41 (2H, m), 1.66 (2H, m), 2.37 (3H, s), 3.50 (2H, t, J = 7.7 Hz), 4.07 (2H, s), 7.01 (1H, d, J = 8.4 Hz), 7.27 (2H, m), 7.49 (1H, d, J = 8.4 Hz), 7.59 (1H, d, J = 2.0 Hz), 7.92 (1H, s), 11.43 (1H, s), 11.78 (1H, s).

IR (Nujol): 1688 cm⁻¹.

Mass (FD): m/e 452 (M).

m.p.: 249 - 250°C.

<Example 62; Production of 3-(2,4-dichlorobenzyl)-2-methyl-5-(1-propanefonylcarbonyl)indole (91)>

According to the method of Example 59, obtained is 3-(2,4-dichlorobenzyl)-2-methyl-5-(1-propanefonylcarbonyl)indole (91) (0.233 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.267 g), N,N'-carbonyldiimidazole (0.195 g), diazabicycloundecene (0.183 g) and 1-propanefonyl (0.148 g).

¹H-NMR (DMSO-d₆, δ ppm): 0.97 (3H, t, J = 7.4 Hz), 1.70 (2H, m), 2.30 (3H, s), 3.46 (2H, t, J = 7.6 Hz), 4.08 (2H, s), 6.93 (1H, d, J = 8.3 Hz), 7.26 (1H, dd, J = 8.3 and 2.1 Hz), 7.35 (1H, d, J = 8.5 Hz), 7.63 (2H, m), 8.04 (1H, s), 11.39 (1H, s), 11.68 (1H, s).

IR (Nujol): 1673 cm⁻¹.

Mass (FD): m/e 438 (M).

m.p.: 161 - 163°C.

<Example 63; Production of 3-(2,4-dichlorobenzyl)-2-methyl-5-(1-octanesulfonylcarbamoyl)indole (92)>

According to the method of Example 59, obtained is 3-(2,4-dichlorobenzyl)-2-methyl-5-(1-octanesulfonylcarbamoyl)indole (92) (0.280 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.300 g), N,N'-carbonyldiimidazole (0.218 g), 1-octanesulfonamide (0.260 g) and diazabicycloundecene (0.205 g).

¹H-NMR (DMSO-d₆, δ ppm): 0.80 (3H, t), 1.12 - 1.27 (8H, m), 1.35 (2H, m), 1.65 (2H, m), 2.30 (3H, s), 3.46 (2H, m), 4.08 (2H, s), 6.92 (1H, d, J = 8.4 Hz), 7.25 (1H, dd, J = 8.4 and 2.1 Hz), 7.34 (1H, d, J = 8.5 Hz), 7.59 - 7.65 (2H, m), 8.03 (1H, s), 11.38 (1H, s), 11.67 (1H, s).

IR (Nujol): 1673 cm⁻¹.

m.p.: 194.5 - 197.5°C.

<Example 64; Production of 5-(benzenesulfonylcarbamoyl)-3-(2,4-dichlorobenzyl)-2-methylindole (93)>

According to the method of Example 59, obtained is 5-(benzenesulfonylcarbamoyl)-3-(2,4-dichlorobenzyl)-2-methylindole (93) (0.235 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.267 g), N,N'-carbonyldiimidazole (0.195 g), benzenesulfonamide (0.189 g) and diazabicycloundecene (0.183 g).

¹H-NMR (DMSO-d₆, δ ppm): 2.28 (3H, s), 4.07 (2H, s), 6.90 (1H, d, J = 8.4 Hz), 7.25 (1H, dd, J = 8.4 and 2.2 Hz), 7.31 (1H, d, J = 8.5 Hz), 7.54 (1H, dd, J = 8.6 and 1.6 Hz), 7.59 - 7.63 (3H, m), 7.68 (1H, m), 7.97 (3H, m), 11.37 (1H, s), 12.19 (1H, brs).

IR (Nujol): 1682 cm⁻¹.

Mass (FD): m/e 472 (M)

m.p.: 244 - 245°C.

<Example 65; Production of 3-(2,4-dichlorobenzyl)-5-(1-hexanesulfonylcarbamoyl)-2-methylindole (94)>

According to the method of Example 59, obtained is 3-(2,4-dichlorobenzyl)-5-(1-hexanesulfonylcarbamoyl)-2-methylindole (94) (0.188 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.267 g), N,N'-carbonyldiimidazole (0.195 g), diazabicycloundecene (0.183 g) and 1-hexanesulfonamide (0.198 g).

$^1\text{H-NMR}$ (DMSO- d_6 , δ ppm): 0.80 (3H, d, $J = 7.0$ Hz), 1.21 (4H, m), 1.36 (2H, m), 1.66 (2H, m), 2.30 (3H, s), 3.47 (2H, t, $J = 7.7$ Hz), 4.08 (2H, s), 6.92 (1H, d, $J = 8.5$ Hz), 7.25 (1H, dd, $J = 8.4$ and 1.2 Hz), 7.35 (1H, d, $J = 8.5$ Hz), 7.62 (2H, m), 8.03 (1H, s), 11.39 (1H, s), 11.67 (1H, s).

IR (Nujol): 1667 cm^{-1} .

Mass (FD): m/e 480 (M)

m.p.: $183 - 185^\circ\text{C}$.

<Example 66; Production of 3-(biphenyl-4-ylmethyl)-2-methyl-5-(1-pentanesulfonylcarbamoyl)indole (95)>

According to the method of Example 59, obtained is 3-(biphenyl-4-ylmethyl)-2-methyl-5-(1-pentanesulfonylcarbamoyl)indole (95) (0.147 g) from 3-(biphenyl-4-ylmethyl)-5-carboxy-2-methylindole (0.355 g), N,N'-carbonyldiimidazole (0.253 g), diazabicycloundecene (0.237 g) and 1-pentanesulfonamide (0.236).

$^1\text{H-NMR}$ (DMSO- d_6 , δ ppm): 0.80 (3H, t, $J = 7.3$ Hz), 1.26 (2H, m), 1.35 (2H, m), 1.68 (2H, m), 2.39 (3H, s), 3.49 (2H, t, $J = 7.7$ Hz), 4.07 (2H, s), 7.29 - 7.33 (4H, m), 7.41 (2H, t, $J = 7.7$ Hz), 7.53 (2H, d, $J = 8.2$ Hz), 7.59 (2H, m), 7.62 (1H, dd, $J = 8.6$ and 1.6 Hz), 8.21 (1H, s), 11.29 (1H, s), 11.73 (1H, s).

IR (Nujol): 1664 cm^{-1} .

Mass (FD): m/e 474 (M)

m.p.: 189 - 190°C.

<Example 67; Production of 3-(2-chlorobenzyl)-2-methyl-5-(1-pentanesulfonylcarbamoyl)indole (96)>

According to the method of Example 59, obtained is 3-(2-chlorobenzyl)-2-methyl-5-(1-pentanesulfonylcarbamoyl)indole (96) (0.115 g) from 5-carboxy-3-(2-chlorobenzyl)-2-methylindole (0.164 g), N,N'-carbonyldiimidazole (0.133 g), diazabicycloundecene (0.125 g) and 1-pentanesulfonamide (0.124).

¹H-NMR (DMSO-d₆, δ ppm): 0.81 (3H, t, J = 7.3 Hz), 1.26 (2H, m), 1.34 (2H, m), 1.66 (2H, m), 2.30 (3H, s), 3.47 (2H, t, J = 7.8 Hz), 4.11 (2H, s), 6.95 (1H, dd, J = 1.6 and 7.6 Hz), 7.15 - 7.21 (2H, m), 7.34 (1H, d, J = 8.5 Hz), 7.45 (1H, dd, J = 1.3 and 7.8 Hz), 7.62 (1H, dd, J = 1.6 and 8.5 Hz), 8.05 (1H, d, J = 1.0 Hz), 11.36 (1H, s), 11.68 (1H, s).

IR (Nujol): 1671 cm⁻¹.

Mass (FD): m/e 432 (M)

m.p.: 175 - 177°C.

<Example 68; Production of 5-(1-butanefulfonylcarbamoyl)-3-(2,4-dichlorobenzoyl)-2-methylindole (97)>

According to the method of Example 59, 5-carboxy-3-(2,4-dichlorobenzoyl)-2-methylindole (0.500 g) is suspended in N,N-dimethylformamide (10 ml), to which is added N,N'-carbonyldiimidazole (0.350 g), and stirred at room temperature for 2 hours. Next, 1-butanefulfonylcarbamoyl (0.296 g) and diazabicycloundecene (0.328 g) are added thereto, and stirred under heat at 100°C for 6 hours. The reaction mixture is concentrated under reduced pressure, and the residue is made acidic with water and 3 M HCl added thereto, and then extracted with ethyl acetate. The organic layer is washed with a saturated saline solution, and dried with anhydrous sodium sulfate. The drying

agent is removed through filtration, and the filtrate is concentrated under reduced pressure. The resulting oily residue is purified through silica gel column chromatography (eluent: chloroform/methanol = 10/1), and recrystallized from ethanol-hexane to obtain a colorless crystal of 5-(1-butanefonylcarbonyl)-3-(2,4-dichlorobenzoyl)-2-methylindole (97) (0.420 g).

$^1\text{H-NMR}$ (DMSO- d_6 , δ ppm): 0.86 (3H, t, $J = 7.3$ Hz), 1.41 (2H, sextet, $J = 7.4$ Hz), 1.66 (2H, quintet, $J = 7.4$ Hz), 2.13 (3H, s), 3.48 (2H, m), 7.47 (1H, d, $J = 8.6$ Hz), 7.48 (1H, d, $J = 8.2$ Hz), 7.58 (1H, dd, $J = 8.0$ and 2.0 Hz), 7.75 (1H, dd, $J = 8.6$ and 1.8 Hz), 7.80 (1H, d, $J = 2.0$ Hz), 8.44 (1H, brs), 12.05 (1H, brs), 12.43 (1H, brs).

IR (Nujol): 1686, 1571 cm^{-1} .

Mass (FD): m/e 466 (M)

m.p.: 241 - 243°C.

<Example 69; Production of 3-(2,4-dichlorobenzyl)-2-methyl-5-(3-methyl-1-butanefonylcarbonyl)indole (98)>

According to the method of Example 59, obtained is 3-(2,4-dichlorobenzyl)-2-methyl-5-(3-methyl-1-butanefonylcarbonyl)indole (98) (0.270 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.290 g), N,N' -carbonyldiimidazole (0.211 g), diazabicycloundecene (0.198 g) and 3-methyl-1-butanefonylamine (0.197 g).

$^1\text{H-NMR}$ (DMSO- d_6 , δ ppm): 0.84 (6H, d, $J = 6.6$ Hz), 1.55 (2H, m), 1.65 (1H, m), 2.30 (2H, s), 3.48 (2H, t, $J = 7.9$ Hz), 4.08 (2H, s), 6.93 (1H, d, $J = 8.4$ Hz), 7.25 (1H, dd, $J = 2.2$ and 8.4 Hz), 7.34 (1H, d, $J = 8.6$ Hz), 7.60 (1H, d, $J = 2.1$ Hz), 7.62 (1H, dd, $J = 1.6$ and 8.6 Hz), 8.02 (1H, s), 11.38 (1H, s), 11.68 (1H, s).

IR (Nujol): 1682 cm^{-1} .

Mass (FD): m/e 466 (M)

m.p.: 167 - 169°C.

<Example 70; Production of 3-(2,4-dichlorobenzyl)-5-(2-methoxyethanesulfonylcarbamoyl)-2-methylindole (99)>

According to the method of Example 59, obtained is 3-(2,4-dichlorobenzyl)-5-(2-methoxyethanesulfonylcarbamoyl)-2-methylindole (99) (0.056 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.290 g), N,N'-carbonyldiimidazole (0.211 g), diazabicycloundecene (0.198 g) and 2-methoxyethanesulfonamide (0.181 g).

¹H-NMR (DMSO-d₆, δ ppm): 2.30 (3H, s), 3.13 (3H, s), 3.59 (4H, m), 4.08 (2H, s), 6.92 (1H, d, J = 8.4 Hz), 7.25 (1H, dd, J = 2.1 and 8.4 Hz), 7.34 (1H, d, J = 8.5 Hz), 7.61 (2H, m), 8.01 (1H, s), 11.37 (1H, s), 11.72 (1H, s).

IR (Nujol): 1671 cm⁻¹.

m.p.: 166 - 168°C.

<Example 71; Production of 3-(4-benzyloxybenzyl)-2-methyl-5-(1-pentanesulfonylcarbamoyl)indole (100)>

According to the method of Example 59, obtained is 3-(4-benzyloxybenzyl)-2-methyl-5-(1-pentanesulfonylcarbamoyl)indole (100) (0.270 g) from 3-(4-benzyloxybenzyl)-5-carboxy-2-methylindole (0.460 g), N,N'-carbonyldiimidazole (0.301 g), diazabicycloundecene (0.283 g) and 1-pentanesulfonamide (0.281 g).

¹H-NMR (DMSO-d₆, δ ppm): 0.81 (3H, t, J = 7.2 Hz), 1.26 (2H, m), 1.35 (2H, m), 1.67 (2H, m), 2.34 (3H, s), 3.43 (2H, t, J = 7.6 Hz), 3.95 (2H, s), 5.01 (2H, s), 6.87 (2H, d, J = 8.1 Hz), 7.11 (2H, d, J = 8.2 Hz), 7.25 - 7.41 (6H, m), 7.61 (1H, d, J = 8.6 Hz), 8.13 (1H, s), 11.17 (1H, s), 11.76 (1H, brs).

IR (Nujol): 1652 cm⁻¹.

Mass (FD): m/e 504 (M).

m.p.: 180 - 184°C.

<Example 72: Production of 3-(2,4-dichlorobenzyl)-5-(1-pentanesulfonylcarbamoyl)-2-propylindole (101)>

According to the method of Example 59, obtained is 3-(2,4-dichlorobenzyl)-5-(1-pentanesulfonylcarbamoyl)-2-propylindole (101) (0.167 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-propylindole (0.185 g), N,N'-carbonyldiimidazole (0.125 g), diazabicycloundecene (0.117 g) and 1-pentanesulfonamide (0.116 g).

$^1\text{H-NMR}$ (DMSO- d_6 , δ ppm): 0.81 (6H, m), 1.25 (2H, m), 1.34 (2H, m), 1.57 (2H, m), 1.66 (2H, m), 2.63 (2H, t, $J=7.5$ Hz), 3.47 (2H, t, $J=7.7$ Hz), 4.09 (2H, s), 6.87 (1H, d, $J=8.4$ Hz), 7.25 (1H, d, $J=8.4$ Hz), 7.36 (1H, d, $J=8.5$ Hz), 7.62 (2H, m), 8.04 (1H, s), 11.38 (1H, s), 11.68 (1H, s).

IR (Nujol): 1682 cm^{-1} .

Mass (FD): m/e 494 (M).

m.p.: $168 - 169^\circ\text{C}$.

<Example 73: Production of 3-(2,4-dichlorobenzyl)-2-ethyl-5-(1-pentanesulfonylcarbamoyl)indole (102)>

According to the method of Example 59, obtained is 3-(2,4-dichlorobenzyl)-2-ethyl-5-(1-pentanesulfonylcarbamoyl)indole (102) (0.255 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-ethylindole (0.370 g), N,N'-carbonyldiimidazole (0.259 g), diazabicycloundecene (0.243 g) and 1-pentanesulfonamide (0.242 g).

$^1\text{H-NMR}$ (DMSO- d_6 , δ ppm): 0.80 (3H, t, $J=7.3$ Hz), 1.14 (3H, t, $J=7.6$ Hz), 1.25 (2H, m), 1.34 (2H, m), 1.66 (2H, m), 2.67 (2H, q, $J=7.6$ Hz), 3.47 (2H, t, $J=7.8$ Hz), 4.09 (2H, s), 6.87 (1H, d, $J=8.4$ Hz), 7.25 (1H, dd, $J=1.9$ and 8.4 Hz), 7.37 (1H, d, $J=8.5$ Hz), 7.63 (2H, m), 8.04 (1H, s), 11.40 (1H, s), 11.68 (1H, s).

IR (Nujol): 1671 cm^{-1} .

Mass (FD): m/e 480 (M).

m.p.: 164.5 - 165.0°C.

<Example 74; Production of 3-(1-bromonaphthalen-2-ylmethyl)-2-methyl-5-(1-pentanesulfonylcarbamoyl)indole (103)>

According to the method of Example 59, obtained is 3-(1-bromonaphthalen-2-ylmethyl)-2-methyl-5-(1-pentanesulfonylcarbamoyl)indole (103) (0.238 g) from 3-(1-bromonaphthalen-2-ylmethyl)-5-carboxy-2-methylindole (0.300 g), N,N'-carbonyldiimidazole (0.247 g), 1-pentanesulfonamide (0.230 g) and diazabicycloundecene (0.232 g).

¹H-NMR (DMSO-d₆, δ ppm): 0.79 (3H, t, J = 7.3 Hz), 1.19 - 1.28 (2H, m), 1.29 - 1.37 (2H, m), 1.61 - 1.69 (2H, m), 2.33 (3H, s), 3.46 (2H, t, J = 7.8 Hz), 4.39 (2H, s), 7.12 (1H, d, J = 8.4 Hz), 7.36 (1H, d, J = 8.5 Hz), 7.56 (1H, t, J = 7.1 Hz), 7.64 (1H, dd, J = 8.5 and 1.6 Hz), 7.68 (1H, t, J = 7.3 Hz), 7.79 (1H, d, J = 8.5 Hz), 7.91 (1H, d, J = 7.9 Hz), 8.10 (1H, s), 8.27 (1H, d, J = 8.6 Hz), 11.40 (1H, s), 11.69 (1H, s).

IR (Nujol): 1674 cm⁻¹.

m.p.: 188.5 - 190.5°C.

<Example 75; Production of 3-((3-chloropyridin-4-yl)methyl)-2-methyl-5-(1-pentanesulfonylcarbamoyl)indole (104)>

According to the method of Example 59, obtained is 3-((3-chloropyridin-4-yl)methyl)-2-methyl-5-(1-pentanesulfonylcarbamoyl)indole (104) (0.263 g) from 5-carboxy-3-((3-chloropyridin-4-yl)methyl)-2-methylindole (0.370 g), N,N'-carbonyldiimidazole (0.309 g), 1-pentanesulfonamide (0.288 g) and diazabicycloundecene (0.290 g).

¹H-NMR (DMSO-d₆, δ ppm): 0.81 (3H, t, J = 7.3 Hz), 1.22 - 1.30 (2H, m), 1.31

- 1.39 (2H, m), 1.63 - 1.71 (2H, m), 2.33 (3H, s), 3.49 (2H, t, $J = 7.8$ Hz), 4.15 (2H, s), 6.92 (1H, d, $J = 5.0$ Hz), 7.37 (1H, d, $J = 8.6$ Hz), 7.64 (1H, d, $J = 8.6$ Hz), 8.06 (1H, s), 8.33 (1H, d, $J = 4.9$ Hz), 8.60 (1H, s), 11.45 (1H, s), 11.69 (1H, s).

IR (Nujol): 1677 cm^{-1} .

m.p.: $217 - 219^{\circ}\text{C}$.

<Example 76; Production of 2-methyl-5-(1-pentanesulfonylcarbamoyl)-3-(4-(2-phenylethenyl)benzyl)indole (105)>

According to the method of Example 59, obtained is 2-methyl-5-(1-pentanesulfonylcarbamoyl)-3-(4-(2-phenylethenyl)benzyl)indole (105) (0.080 g) from 5-carboxy-2-methyl-3-(4-(2-phenylethenyl)benzyl)indole (0.120 g), N,N'-carbonyldiimidazole (0.106 g), 1-pentanesulfonamide (0.099 g) and diazabicycloundecene (0.100 g).

$^1\text{H-NMR}$ (CD_3OD , δ ppm): 0.79 (3H, t, $J = 7.3$ Hz), 1.20 - 1.38 (4H, m), 1.71 (2H, m), 2.32 (3H, s), 3.42 (2H, m), 4.02 (2H, s), 7.01 (2H, d, $J = 3.8$ Hz), 7.09 - 7.14 (3H, m), 7.20 - 7.27 (3H, m), 7.34 (2H, d, $J = 8.1$ Hz), 7.41 (2H, d, $J = 7.3$ Hz), 7.53 (1H, d, $J = 10.2$ Hz), 7.95 (1H, s).

IR (Nujol): 1666 cm^{-1} .

m.p.: $216 - 219^{\circ}\text{C}$.

<Example 77; Production of 3-(2,4-dichlorobenzyl)-5-(ethanesulfonylcarbamoyl)-2-propylindole (106)>

According to the method of Example 59, obtained is 3-(2,4-dichlorobenzyl)-5-(ethanesulfonylcarbamoyl)-2-propylindole (106) (0.130 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-propylindole (0.154 g), N,N'-carbonyldiimidazole (0.138 g), diazabicycloundecene (0.129 g) and ethanesulfonamide (0.0928 g).

$^1\text{H-NMR}$ (DMSO- d_6 , δ ppm): 0.82 (3H, t, $J = 7.3$ Hz), 1.22 (3H, t, $J = 7.3$ Hz), 1.57 (2H, m), 2.62 (2H, t, $J = 7.5$ Hz), 3.48 (2H, q, $J = 7.3$ Hz), 4.10 (2H, s), 6.88 (1H, d, $J = 8.4$ Hz), 7.25 (1H, dd, $J = 8.4$ and 2.1 Hz), 7.37 (1H, d, $J = 7.4$ Hz), 7.61 (1H, d, $J = 2.1$ Hz), 7.65 (1H, dd, $J = 8.5$ and 1.6 Hz), 8.05 (1H, brs), 11.38 (1H, s), 11.66 (1H, s).

IR (Nujol): 1683 cm^{-1} .

m.p.: $204.5 - 205.5^\circ\text{C}$.

<Example 78; Production of 3-(2,4-dichlorobenzyl)-2-methyl-5-(2-thiophenesulfonylcarbamoyl)indole (107)>

According to the method of Example 59, obtained is 3-(2,4-dichlorobenzyl)-2-methyl-5-(2-thiophenesulfonylcarbamoyl)indole (107) (0.226 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.334 g), N,N' -carbonyldiimidazole (0.275 g), diazabicycloundecene (0.258 g) and 2-thiophenesulfonamide (0.277 g).

$^1\text{H-NMR}$ (DMSO- d_6 , δ ppm): 2.28 (3H, s), 4.07 (2H, s), 6.91 (1H, d, $J = 8.4$ Hz), 7.18 (1H, t, $J = 4.2$ Hz), 7.25 (1H, dd, $J = 2.2$ and 8.4 Hz), 7.32 (1H, d, $J = 8.5$ Hz), 7.58 (1H, dd, $J = 1.6$ and 8.6 Hz), 7.61 (1H, d, $J = 2.1$ Hz), 7.82 (1H, dd, $J = 1.0$ and 3.8 Hz), 8.00 (2H, m), 11.39 (1H, brs), 12.31 (1H, brs).

IR (Nujol): 1690 cm^{-1} .

m.p.: $221 - 222^\circ\text{C}$.

<Example 79; Production of 3-(2,4-dichlorobenzyl)-5-((4-methoxybenzene)sulfonylcarbamoyl)-2-methylindole (108)>

According to the method of Example 59, obtained is 3-(2,4-dichlorobenzyl)-5-((4-methoxybenzene)sulfonylcarbamoyl)-2-methylindole (108) (0.260 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.300 g), N,N' -carbonyldiimidazole (0.185 g), 4-methoxybenzenesulfonamide (0.214 g) and diazabicycloundecene (0.174

g).

$^1\text{H-NMR}$ (DMSO- d_6 , δ ppm): 2.29 (3H, s), 3.83 (3H, s), 4.07 (2H, s), 6.92 (1H, d, $J = 8.4$ Hz), 7.13 (2H, d, $J = 9.0$ Hz), 7.26 (1H, dd, $J = 8.4$ and 2.3 Hz), 7.31 (1H, d, $J = 8.6$ Hz), 7.54 (1H, dd, $J = 8.5$ and 1.8 Hz), 7.62 (1H, d, $J = 2.2$ Hz), 7.91 (2H, d, $J = 8.9$ Hz), 7.97 (1H, s), 11.37 (1H, s), 12.03 (1H, s).

IR (Nujol): 1684 cm^{-1} .

m.p.: $106.5 - 109^\circ\text{C}$.

<Example 80; Production of 5-(benzenesulfonylcarbamoyl)-3-(2,4-dichlorobenzyl)-2-ethylindole (109)>

According to the method of Example 59, obtained is 5-(benzenesulfonylcarbamoyl)-3-(2,4-dichlorobenzyl)-2-ethylindole (109) (0.095 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-ethylindole (0.313 g), N,N'-carbonyldiimidazole (0.212 g), diazabicycloundecene (0.206 g) and benzenesulfonamide (0.219 g).

$^1\text{H-NMR}$ (DMSO- d_6 , δ ppm): 1.13 (3H, t, $J = 7.5$ Hz), 2.65 (2H, q, $J = 7.5$ Hz), 4.08 (2H, s), 6.85 (1H, d, $J = 8.4$ Hz), 7.24 (1H, dd, $J = 8.4$ and 2.1 Hz), 7.33 (1H, d, $J = 8.6$ Hz), 7.54 - 7.64 (4H, m), 7.68 (1H, t), 7.95 - 8.00 (3H, m), 11.38 (1H, s), 12.18 (1H, s).

IR (Nujol): 1696 cm^{-1} .

m.p.: $229 - 231^\circ\text{C}$.

<Example 81; Production of 3-((4-chloroisoquinolin-3-yl)methyl)-2-methyl-5-(1-pentanesulfonylcarbamoyl)indole (110)>

According to the method of Example 59, obtained is 3-((4-chloroisoquinolin-3-yl)methyl)-2-methyl-5-(1-pentanesulfonylcarbamoyl)indole (110) (0.020 g) from 5-carboxy-3-((4-chloroisoquinolin-3-yl)methyl)-2-methylindole (0.112 g), N,N'-carbonyldiimidazole (0.078 g), diazabicycloundecene (0.073 g) and 1-

pentanesulfonamide (0.072 g).

$^1\text{H-NMR}$ (DMSO- d_6 , δ ppm): 0.79 (3H, t, $J = 7.3$ Hz), 1.25 (2H, m), 1.34 (2H, m), 1.66 (2H, m), 2.41 (3H, s), 3.43 (2H, t, $J = 6.5$ Hz), 4.49 (2H, s), 7.26 (1H, d, $J = 8.5$ Hz), 7.56 (1H, d, $J = 8.5$ Hz), 7.72 (1H, t, $J = 7.6$ Hz), 7.91 (1H, t, $J = 7.7$ Hz), 8.16 (2H, m), 8.26 (1H, s), 9.22 (1H, s), 11.22 (1H, s), 11.71 (1H, brs).

IR (Nujol): 1673 cm^{-1} .

m.p.: $196 - 198^\circ\text{C}$.

<Example 82; Production of 3-((4-bromoisoquinolin-3-yl)methyl)-2-methyl-5-(1-pentanesulfonylcarbamoyl)indole (111)>

According to the method of Example 59, obtained is 3-((4-bromoisoquinolin-3-yl)methyl)-2-methyl-5-(1-pentanesulfonylcarbamoyl)indole (111) (0.063 g) from 3-((4-bromoisoquinolin-3-yl)methyl)-5-carboxy-2-methylindole (0.121 g), N,N' -carbonyldiimidazole (0.099 g), diazabicycloundecene (0.093 g) and 1-pentanesulfonamide (0.093 g).

$^1\text{H-NMR}$ (DMSO- d_6 , ppm): 0.97 (3H, t, $J = 7.3$ Hz), 1.24 (2H, m), 1.34 (2H, m), 1.66 (2H, m), 2.39 (3H, s), 3.43 (2H, m), 4.54 (2H, s), 7.26 (1H, d, $J = 8.5$ Hz), 7.56 (1H, dd, $J = 1.6$ and 8.5 Hz), 7.72 (1H, t, $J = 7.7$ Hz), 7.91 (1H, m), 8.14 (2H, m), 8.23 (1H, s), 9.22 (1H, s), 11.22 (1H, s), 11.69 (1H, brs).

IR (Nujol): 1676 cm^{-1} .

m.p.: $171 - 175^\circ\text{C}$.

<Example 83; Production of 3-(2,4-dichlorobenzyl)-2-methyl-5-(1-pent-1-enesulfonylcarbamoyl)indole (112)>

According to the method of Example 59, obtained is 3-(2,4-dichlorobenzyl)-2-methyl-5-(1-pent-1-enesulfonylcarbamoyl)indole (112) (0.300 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.325 g), N,N' -carbonyldiimidazole (0.315 g), 1-

pent-1-enesulfonamide (0.290 g) and diazabicycloundecene (0.290 g).

$^1\text{H-NMR}$ (CDCl_3 , δ ppm): 0.89 (3H, d, $J = 7.4$ Hz), 1.46 (2H, m), 2.19 (2H, m), 2.32 (3H, s), 4.05 (2H, s), 6.62 (1H, d, $J = 15.2$ Hz), 6.75 (1H, d, $J = 8.4$ Hz), 6.97 (1H, dd, $J = 8.3$ and 2.1 Hz), 7.03 (1H, m), 7.30 (1H, d, $J = 8.5$ Hz), 7.37 (1H, d, $J = 2.1$ Hz), 7.57 (1H, dd, $J = 8.5$ and 1.6 Hz), 7.87 (1H, s), 8.62 (1H, s).

IR (Nujol): 1667 cm^{-1} .

<Example 84; Production of 3-(2,4-dichlorobenzyl)-2-methyl-5-(trifluoromethanesulfonylcarbamoyl)indole (113)>

According to the method of Example 59, obtained is 3-(2,4-dichlorobenzyl)-2-methyl-5-(trifluoromethanesulfonylcarbamoyl)indole (113) (0.088 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.414 g), N,N' -carbonyldiimidazole (0.40 g), diazabicycloundecene (0.378 g) and trifluoromethanesulfonamide (0.307 g).

$^1\text{H-NMR}$ (DMSO-d_6 , δ ppm): 2.31 (3H, s), 4.05 (2H, s), 6.96 (1H, d, $J = 8.4$ Hz), 7.24 (2H, m), 7.58 (1H, d, $J = 2.2$ Hz), 7.65 (1H, dd, $J = 1.3$ and 8.5 Hz), 7.89 (1H, s), 11.12 (1H, s).

IR (Nujol): 1722 cm^{-1} .

Mass (FD): m/e 464 (M).

m.p.: $201 - 205^\circ\text{C}$.

<Example 85; Production of 3-(2,4-dichlorobenzyl)-5-(2,2-dimethylpropanesulfonylcarbamoyl)-2-methylindole (114)>

According to the method of Example 59, obtained is 3-(2,4-dichlorobenzyl)-5-(2,2-dimethylpropanesulfonylcarbamoyl)-2-methylindole (114) (0.287 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.368 g), N,N' -carbonyldiimidazole (0.268 g), diazabicycloundecene (0.251 g) and 2,2-dimethylpropanesulfonamide (0.250 g).

$^1\text{H-NMR}$ (DMSO- d_6 , δ ppm): 1.09 (9H, s), 2.30 (3H, s), 3.48 (2H, s), 4.08 (2H, s), 6.93 (1H, d, $J = 8.4$ Hz), 7.26 (1H, dd, $J = 1.9$ and 8.3 Hz), 7.34 (1H, d, $J = 8.6$ Hz), 7.61 (1H, d, $J = 2.0$ Hz), 7.62 (1H, d, $J = 8.5$ Hz), 8.02 (1H, s), 11.38 (1H, s), 11.70 (1H, brs).

IR (Nujol): 1656 cm^{-1} .

m.p.: $134 - 138^\circ\text{C}$.

<Example 86; Production of 3-(2,4-dichlorobenzyl)-2-methyl-5-(8-quinolinesulfonylcarbamoyl)indole (115)>

According to the method of Example 59, obtained is 3-(2,4-dichlorobenzyl)-2-methyl-5-(8-quinolinesulfonylcarbamoyl)indole (115) (0.170 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.200 g), N,N' -carbonyldiimidazole (0.198 g), 8-quinolinesulfonamide (0.249 g) and diazabicycloundecene (0.182 g).

$^1\text{H-NMR}$ (DMSO- d_6 , δ ppm): 2.27 (3H, s), 4.02 (2H, s), 6.81 (1H, d, $J = 8.4$ Hz), 7.21 - 7.27 (2H, m), 7.47 (1H, d, $J = 9.6$ Hz), 7.55 (1H, dd, $J = 8.2$ and 4.2 Hz), 7.66 (1H, d, $J = 2.0$ Hz), 7.81 (1H, t, $J = 7.8$ Hz), 8.06 (1H, s), 8.32 (1H, d, $J = 8.0$ Hz), 8.50 (2H, d, $J = 7.6$ Hz), 8.76 (1H, d), 11.32 (1H, s).

IR (Nujol): 1678 cm^{-1} .

m.p.: $255 - 256^\circ\text{C}$.

<Example 87; Production of 3-(2,4-dichlorobenzyl)-2-methyl-5-((2-phenylethane)sulfonylcarbamoyl)indole (116)>

According to the method of Example 59, obtained is 3-(2,4-dichlorobenzyl)-2-methyl-5-((2-phenylethane)sulfonylcarbamoyl)indole (116) (0.050 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.145 g), N,N' -carbonyldiimidazole (0.100 g), 8-quinolinesulfonamide (0.114 g) and diazabicycloundecene (0.094 g).

$^1\text{H-NMR}$ (DMSO- d_6 , δ ppm): 2.30 (3H, s), 3.01 (2H, m), 3.80 (2H, m), 4.09

(2H, s), 6.93 (1H, d, $J = 8.4$ Hz), 7.15 (1H, m), 7.21 - 7.28 (5H, m), 7.34 (1H, d, $J = 8.5$ Hz), 7.60 - 7.64 (2H, m), 8.01 (1H, s), 11.39 (1H, s), 11.77 (1H, s).

IR (Nujol): 1674 cm^{-1} .

m.p.: $212 - 213^{\circ}\text{C}$.

<Example 88: Production of 3-(2,4-dichlorobenzyl)-2-methyl-5-(-toluenesulfonylcarbamoyl)indole (117)>

According to the method of Example 59, obtained is 3-(2,4-dichlorobenzyl)-2-methyl-5-(-toluenesulfonylcarbamoyl)indole (117) (0.217 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.334 g), N,N'-carbonyldiimidazole (0.243 g), diazabicycloundecene (0.228 g) and -toluenesulfonamide (0.257 g).

[Physical Properties of Compound (117)]

^1H -NMR (DMSO- d_6 , ppm): 2.30 (3H, s), 4.05 (2H, s), 4.81 (2H, s), 6.90 (1H, d, $J = 8.4$ Hz), 7.26 (1H, dd, $J = 2.1$ and 8.4 Hz), 7.29 (5H, m), 7.35 (1H, d, $J = 8.5$ Hz), 7.61 (1H, d, $J = 2.1$ Hz), 7.63 (1H, dd, $J = 1.7$ and 8.6 Hz), 7.97 (1H, d, $J = 1.0$ Hz), 11.46 (1H, s), 11.62 (1H, s).

IR (Nujol): 1690 cm^{-1} .

m.p.: $224 - 225^{\circ}\text{C}$.

<Example 89: Production of 5-cyclohexanesulfonylcarbamoyl-3-(2,4-dichlorobenzyl)-2-methylindole (118)>

According to the method of Example 59, obtained is 5-cyclohexanesulfonylcarbamoyl-3-(2,4-dichlorobenzyl)-2-methylindole (118) (0.032 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.217 g), N,N'-carbonyldiimidazole (0.126 g), diazabicycloundecene (0.119 g) and cyclohexanesulfonamide (0.127 g).

^1H -NMR (DMSO- d_6 , δ ppm): 1.14 (1H, m), 1.26 (2H, m), 1.46 (2H, m), 1.60

(1H, m), 1.79 (2H, d, $J = 11.6$ Hz), 2.00 (2H, d, $J = 10.4$ Hz), 2.29 (3H, s), 3.57

(1H, m), 4.08 (2H, s), 6.93 (1H, d, $J = 8.3$ Hz), 7.26 (1H, d, $J = 8.2$ Hz), 7.34

(1H, d, $J = 8.4$ Hz), 7.62 (2H, m), 8.03 (1H, s), 11.39 (1H, s), 11.60 (1H, brs).

IR (Nujol): 1657 cm^{-1} .

m.p.: $176 - 181^{\circ}\text{C}$.

<Example 90; Production of 5-(3-chloro-1-propanesulfonylcarbamoyl)-3-(2,4-dichlorobenzyl)-2-methylindole (119) and 3-(2,4-dichlorobenzyl)-2-methyl-5-(propanesultam-1-ylcarbonyl)indole (120)>

A mixture of 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.334 g), N,N'-carbonyldiimidazole (0.243 g) and N,N-dimethylformamide (5 ml) is stirred at room temperature for 40 minutes. Then, (3-chloro-1-propane)sulfonamide (0.236 g) and diazabicycloundecene (0.228 g) are added thereto, and stirred at room temperature for 14 days. This is made acidic with 0.5 M HCl (8 ml), and the solid formed is taken out through filtration. This solid is dissolved in ethyl acetate (60 ml), to which is added a saline solution (100 ml) for liquid-liquid separation. The organic layer is dried and concentrated, and the resulting residue is separated and purified through silica gel column chromatography (methanol/chloroform = 3/97). The resulting two components are separately recrystallized and purified from diethyl ether/hexane to obtain 5-(3-chloro-1-propanesulfonylcarbamoyl)-3-(2,4-dichlorobenzyl)-2-methylindole (119) (0.160 g) and 3-(2,4-dichlorobenzyl)-2-methyl-5-(propanesultam-1-ylcarbonyl)indole (120) (0.091 g).

$^1\text{H-NMR}$ (DMSO- d_6 , δ ppm): 2.13 (2H, m), 2.30 (3H, s), 3.63 (2H, t, $J = 7.5$ Hz), 3.75 (2H, t, $J = 6.5$ Hz), 4.09 (2H, s), 6.93 (1H, d, $J = 8.4$ Hz), 7.26 (1H, dd, $J = 2.1$ and 8.4 Hz), 7.35 (1H, d, $J = 8.5$ Hz), 7.61 (1H, d, $J = 2.1$ Hz), 7.63 (1H, dd, $J = 1.6$ and 8.6 Hz), 8.03 (1H, s), 11.40 (1H, s), 11.80 (1H, brs).

IR (Nujol): 1681 cm^{-1} .

m.p.: 95.5 - 96.2°C.

[Physical Properties of Compound (120)]

^1H -NMR (DMSO- d_6 , δ ppm): 2.29 (2H, m), 2.33 (3H, s), 3.49 (2H, t, $J = 7.0$ Hz), 3.85 (2H, t, $J = 6.6$ Hz), 4.06 (2H, s), 7.02 (1H, d, $J = 8.4$ Hz), 7.25 (1H, m), 7.33 (1H, d, $J = 8.4$ Hz), 7.39 (1H, dd, $J = 1.4$ and 8.4 Hz), 7.58 (1H, m), 7.72 (1H, s), 11.33 (1H, s).

IR (Nujol): 1658 cm^{-1} .

m.p.: 197 -198°C.

<Example 91; Production of 6-(1-butanefonylcarbamoil)-2-(2,4-dichlorobenzyl)-3-methylindole (121)>

From 6-carboxy-2-(2,4-dichlorobenzyl)-3-methylindole (0.550 g), N,N'-carbonyldiimidazole (0.400 g), 1-butanefonamide (0.340 g) and diazabicycloundecene (0.376 g), obtained is 6-(1-butanefonylcarbamoil)-2-(2,4-dichlorobenzyl)-3-methylindole (121) (0.250 g).

^1H -NMR (DMSO- d_6 , δ ppm): 0.86 (3H, t, $J = 7.3$ Hz), 1.40 (2H, sextet, $J = 7.4$ Hz), 1.66 (2H, quint, $J = 7.4$ Hz), 2.18 (3H, s), 3.50 (2H, m), 4.19 (2H, s), 7.09 (1H, d, $J = 8.4$ Hz), 7.36 (1H, dd, $J = 2.2$ and 8.4 Hz), 7.51 (1H, d, $J = 8.4$ Hz), 7.57 (1H, dd, $J = 1.5$ and 8.4 Hz), 7.62 (1H, d, $J = 2.2$ Hz), 7.91 (1H, d, $J = 1.5$ Hz), 11.16 (1H, brs), 11.77 (1H, brs).

IR (Nujol): 1666 cm^{-1} .

Mass (FD): m/e 452 (M).

m.p.: 215.5 -216.1°C.

<Example 92; Production of 1-(2,4-dichlorobenzyl)-3-methyl-6-(1-pentanesufonylcarbamoil)indole (122)>

According to the method of Example 59, obtained is a colorless crystal of 1-(2,4-dichlorobenzyl)-3-methyl-6-(1-pentanesulfonylcarbamoyl)indole (122) (0.370 g) from 6-carboxy-1-(2,4-dichlorobenzyl)-3-methylindole (0.46 g), N,N'-carbonyldiimidazole (0.335 g), 1-pentanesulfonamide (0.312 g) and diazabicycloundecene (0.314 g).

$^1\text{H-NMR}$ (DMSO- d_6 , δ ppm): 0.80 (3H, t, $J = 7.3$ Hz), 1.25 (2H, sextet, $J = 7.6$ Hz), 1.34 (2H, quint, $J = 7.5$ Hz), 1.67 (2H, quint, $J = 7.5$ Hz), 2.28 (3H, s), 3.49 (2H, m), 5.50 (2H, s), 6.55 (1H, d, $J = 8.4$ Hz), 7.32 (1H, dd, $J = 2.0$ and 8.4 Hz), 7.42 (1H, s), 7.64 (2H, s), 7.70 (1H, d, $J = 2.0$ Hz), 8.10 (1H, s), 11.77 (1H, brs).

IR (Nujol): 1686 cm^{-1} .

m.p.: $116 - 117.5^\circ\text{C}$.

<Example 93; Production of 3-(2,4-dichlorobenzyl)-2-methyl-5-((4-methylbenzene)sulfonylcarbamoyl)indole (123)>

According to the method of Example 59, obtained is 3-(2,4-dichlorobenzyl)-2-methyl-5-((4-methylbenzene)sulfonylcarbamoyl)indole (123) (0.356 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.334 g), N,N'-carbonyldiimidazole (0.243 g), (4-methylbenzene)sulfonamide (0.257 g) and diazabicycloundecene (0.228 g).

$^1\text{H-NMR}$ (DMSO- d_6 , δ ppm): 2.28 (3H, s), 2.37 (3H, s), 4.06 (2H, s), 6.90 (1H, d, $J = 8.4$ Hz), 7.25 (1H, dd, $J = 8.4$ and 2.2 Hz), 7.30 (1H, d, $J = 8.6$ Hz), 7.40 (2H, d, $J = 8.2$ Hz), 7.53 (1H, dd, $J = 8.7$ and 2.1 Hz), 7.60 (1H, d, $J = 2.2$ Hz), 7.85 (2H, d, $J = 8.3$ Hz), 7.96 (1H, s), 11.36 (1H, s), 12.09 (1H, s).

IR (Nujol): 1688 cm^{-1} .

m.p.: $243 - 246^\circ\text{C}$.

<Example 94; Production of 3-(2,4-dichlorobenzyl)-2-methyl-5-((4-nitrobenzene)sulfonylcarbamoyl)indole (124)>

According to the method of Example 59, obtained is 3-(2,4-dichlorobenzyl)-2-methyl-5-((4-nitrobenzene)sulfonylcarbamoyl)indole (124) (0.350 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.334 g), N,N'-carbonyldiimidazole (0.243 g), (4-nitrobenzene)sulfonamide (0.303 g) and diazabicycloundecene (0.228 g).

$^1\text{H-NMR}$ (DMSO- d_6 , δ ppm): 2.28 (3H, s), 4.07 (2H, s), 6.89 (1H, d, $J = 8.4$ Hz), 7.25 (1H, dd, $J = 8.4$ and 2.1 Hz), 7.32 (1H, d, $J = 8.5$ Hz), 7.55 (1H, d, $J = 8.5$ Hz), 7.60 (1H, d, $J = 2.0$ Hz), 7.97 (1H, s), 8.21 (2H, d, $J = 8.8$ Hz), 8.43 (2H, d, $J = 8.9$ Hz), 11.41 (1H, s), 12.50 (1H, brs).

IR (Nujol): 1675 cm^{-1} .

m.p.: $216 - 221^\circ\text{C}$.

<Example 95; Production of 5-((4-chlorobenzene)sulfonylcarbamoyl)-3-(2,4-dichlorobenzyl)-2-methylindole (125)>

According to the method of Example 59, obtained is 5-((4-chlorobenzene)sulfonylcarbamoyl)-3-(2,4-dichlorobenzyl)-2-methylindole (125) (0.195 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.334 g), N,N'-carbonyldiimidazole (0.243 g), (4-chlorobenzene)sulfonamide (0.287 g) and diazabicycloundecene (0.228 g).

$^1\text{H-NMR}$ (DMSO- d_6 , δ ppm): 2.28 (3H, s), 4.07 (2H, s), 6.90 (1H, d, $J = 8.5$ Hz), 7.25 (1H, dd, $J = 8.4$ and 2.3 Hz), 7.32 (1H, d, $J = 8.6$ Hz), 7.54 (1H, dd, $J = 8.7$ and 1.5 Hz), 7.61 (1H, d, $J = 2.0$ Hz), 7.69 (2H, d, $J = 8.6$ Hz), 7.97 (3H, m), 11.39 (1H, s), 12.27 (1H, brs).

IR (Nujol): 1659 cm^{-1} .

m.p.: $133 - 136^\circ\text{C}$.

<Example 96; Production of 5-((3-chlorobenzene)sulfonylcarbamoyl)-3-(2,4-dichlorobenzyl)-2-methylindole (126)>

According to the method of Example 59, obtained is 5-((3-chlorobenzene)sulfonylcarbamoyl)-3-(2,4-dichlorobenzyl)-2-methylindole (126) (0.107 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.334 g), N,N'-carbonyldiimidazole (0.243 g), (3-chlorobenzene)sulfonamide (0.287 g) and diazabicycloundecene (0.228 g).

$^1\text{H-NMR}$ (DMSO- d_6 , δ ppm): 2.28 (3H, s), 4.07 (2H, s), 6.90 (1H, d, $J = 8.5$ Hz), 7.25 (1H, dd, $J = 8.4$ and 2.2 Hz), 7.32 (1H, d, $J = 8.5$ Hz), 7.55 (1H, m), 7.60 (1H, d, $J = 2.1$ Hz), 7.66 (1H, t, $J = 8.0$ Hz), 7.79 (1H, d, $J = 7.8$ Hz), 7.92 (1H, d, $J = 7.7$ Hz), 7.95 (1H, s), 7.98 (1H, s), 11.40 (1H, s), 12.31 (1H, brs).

IR (Nujol): 1687 cm^{-1} .

m.p.: $206 - 207^\circ\text{C}$.

<Example 97; Production of 5-((2-chlorobenzene)sulfonylcarbamoyl)-3-(2,4-dichlorobenzyl)-2-methylindole (127)>

According to the method of Example 59, obtained is 5-((2-chlorobenzene)sulfonylcarbamoyl)-3-(2,4-dichlorobenzyl)-2-methylindole (127) (0.240 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.334 g), N,N'-carbonyldiimidazole (0.243 g), (2-chlorobenzene)sulfonamide (0.287 g) and diazabicycloundecene (0.228 g).

$^1\text{H-NMR}$ (DMSO- d_6 , δ ppm): 2.29 (3H, s), 4.07 (2H, s), 6.94 (1H, d, $J = 8.4$ Hz), 7.26 (1H, dd, $J = 2.0$ and 8.3 Hz), 7.31 (1H, d, $J = 8.6$ Hz), 7.56 (1H, d, $J = 8.5$ Hz), 7.61 (3H, m), 7.68 (1H, m), 8.06 (1H, s), 8.15 (1H, d, $J = 7.8$ Hz), 11.38 (1H, s), 12.54 (1H, brs).

IR (Nujol): 1690 cm^{-1} .

m.p.: $224 - 226^\circ\text{C}$.

<Example 98; Production of 3-(2,4-dichlorobenzyl)-5-((4-

fluorobenzene)sulfonylcarbamoyl)-2-methylindole (128)>

According to the method of Example 59, obtained is 3-(2,4-dichlorobenzyl)-5-((4-fluorobenzene)sulfonylcarbamoyl)-2-methylindole (128) (0.130 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.334 g), N,N'-carbonyldiimidazole (0.243 g), (4-fluorobenzene)sulfonamide (0.263 g) and diazabicycloundecene (0.228 g).

$^1\text{H-NMR}$ (DMSO- d_6 , δ ppm): 2.28 (3H, s), 4.07 (2H, s), 6.90 (1H, d, $J = 8.4$ Hz), 7.25 (1H, dd, $J = 1.9$ and 8.4 Hz), 7.32 (1H, d, $J = 8.6$ Hz), 7.45 (2H, t, $J = 8.7$ Hz), 7.55 (1H, d, $J = 8.5$ Hz), 7.60 (1H, d, $J = 2.0$ Hz), 7.97 (1H, s), 8.03 (1H, d, $J = 8.6$ Hz), 8.04 (1H, d, $J = 8.8$ Hz), 11.38 (1H, s), 12.22 (1H, brs).

IR (Nujol): 1685 cm^{-1} .

m.p.: $222 - 225^\circ\text{C}$.

<Example 99; Production of 3-(2,4-dichlorobenzyl)-2-methyl-5-((2-naphthalene)sulfonylcarbamoyl)indole (129)>

According to the method of Example 59, obtained is 3-(2,4-dichlorobenzyl)-2-methyl-5-(2-naphthalenesulfonylcarbamoyl)indole (129) (0.132 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.334 g), N,N'-carbonyldiimidazole (0.243 g), 2-naphthalenesulfonamide (0.331 g) and diazabicycloundecene (0.228 g).

$^1\text{H-NMR}$ (DMSO- d_6 , δ ppm): 2.27 (3H, s), 4.06 (2H, s), 6.90 (1H, d, $J = 8.4$ Hz), 7.24 (1H, dd, $J = 8.4$ and 1.6 Hz), 7.30 (1H, d, $J = 8.5$ Hz), 7.53 (1H, d, $J = 8.5$ Hz), 7.59 (1H, d, $J = 2.2$ Hz), 7.68 (1H, t, $J = 8.1$ Hz), 7.73 (1H, t, $J = 7.9$ Hz), 7.94 (1H, dd, $J = 8.7$ and 1.7 Hz), 7.98 (1H, s), 8.04 (1H, d, $J = 8.1$ Hz), 8.12 (1H, d, $J = 8.7$ Hz), 8.22 (1H, d, $J = 8.1$ Hz), 8.65 (1H, s), 11.36 (1H, s), 12.27 (1H, brs).

IR (Nujol): 1688 cm^{-1} .

m.p.: $219 - 224^\circ\text{C}$.

<Example 100; Production of 3-(2,4-dichlorobenzyl)-2-methyl-5-((1-naphthalene)sulfonylcarbamoyl)indole (130)>

According to the method of Example 59, obtained is 3-(2,4-dichlorobenzyl)-2-methyl-5-((1-naphthalenesulfonylcarbamoyl)indole (130) (0.191 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.334 g), N,N'-carbonyldiimidazole (0.243 g), 1-naphthalenesulfonamide (0.331 g) and diazabicycloundecene (0.228 g).

$^1\text{H-NMR}$ (DMSO- d_6 , δ ppm): 2.27 (3H, s), 4.06 (2H, s), 6.85 (1H, d, $J = 8.4$ Hz), 7.21 (1H, dd, $J = 2.2$ and 8.3 Hz), 7.27 (1H, d, $J = 8.6$ Hz), 7.48 (1H, dd, $J = 1.7$ and 8.6 Hz), 7.60 (2H, m), 7.63 (1H, d, $J = 2.2$ Hz), 7.72 (1H, t, $J = 7.8$ Hz), 7.94 (1H, s), 8.08 (1H, m), 8.28 (1H, d, $J = 8.3$ Hz), 8.36 (1H, d, $J = 7.4$ Hz), 8.69 (1H, m), 11.35 (1H, s), 12.50 (1H, brs).

IR (Nujol): 1693 cm^{-1} .

m.p.: $221 - 223^\circ\text{C}$.

<Example 101; Production of 3-(2,4-dichlorobenzyl)-2-methyl-5-((2-methylbenzene)sulfonylcarbamoyl)indole (131)>

According to the method of Example 59, obtained is 3-(2,4-dichlorobenzyl)-2-methyl-5-((2-methylbenzene)sulfonylcarbamoyl)indole (131) (0.150 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.334 g), N,N'-carbonyldiimidazole (0.243 g), (2-methylbenzene)sulfonamide (0.257 g) and diazabicycloundecene (0.228 g).

$^1\text{H-NMR}$ (DMSO- d_6 , δ ppm): 2.29 (3H, s), 2.57 (3H, s), 4.07 (2H, s), 6.94 (1H, d, $J = 8.4$ Hz), 7.25 (1H, dd, $J = 2.1$ and 8.4 Hz), 7.31 (1H, t, $J = 7.6$ Hz), 7.36 (1H, d, $J = 7.4$ Hz), 7.43 (1H, t, $J = 7.6$ Hz), 7.55 (2H, m), 7.60 (1H, d, $J = 2.1$ Hz), 8.01 (2H, m), 11.37 (1H, s), 12.28 (1H, brs).

IR (Nujol): 1689 cm^{-1} .

m.p.: $194 - 196^\circ\text{C}$.

<Example 102; Production of 3-(2,4-dichlorobenzyl)-5-(2,5-dimethylbenzene)sulfonylcarbamoyl-2-methylindole (132)>

According to the method of Example 59, obtained is 3-(2,4-dichlorobenzyl)-5-(2,5-dimethylbenzene)sulfonylcarbamoyl-2-methylindole (132) (0.285 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.334 g), N,N'-carbonyldiimidazole (0.243 g), (2,5-dimethylbenzene)sulfonamide (0.278 g) and diazabicycloundecene (0.228 g).

¹H-NMR (DMSO-d₆, δ ppm): 2.29 (3H, s), 2.35 (3H, s), 2.51 (3H, s), 4.07 (2H, s), 6.93 (1H, d, J = 8.4 Hz), 7.25 (2H, m), 7.31 (1H, d, J = 8.5 Hz), 7.36 (1H, d, J = 7.8 Hz), 7.55 (1H, dd, J = 1.5 and 8.5 Hz), 7.60 (1H, d, J = 2.1 Hz), 7.82 (1H, s), 8.01 (1H, s), 11.37 (1H, s), 12.24 (1H, s).

IR (Nujol): 1682 cm⁻¹.

m.p.: 228 - 238°C.

<Example 103; Production of 5-(4-bromobenzene)sulfonylcarbamoyl-3-(2,4-dichlorobenzyl)-2-methylindole (133)>

According to the method of Example 59, obtained is 5-(4-bromobenzene)sulfonylcarbamoyl-3-(2,4-dichlorobenzyl)-2-methylindole (133) (0.295 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.334 g), N,N'-carbonyldiimidazole (0.243 g), (4-bromobenzene)sulfonamide (0.354 g) and diazabicycloundecene (0.228 g).

¹H-NMR (DMSO-d₆, δ ppm): 2.28 (3H, s), 4.07 (2H, s), 6.90 (1H, d, J = 8.4 Hz), 7.25 (1H, dd, J = 1.7 and 8.3 Hz), 7.32 (1H, d, J = 8.5 Hz), 7.54 (1H, d, J = 8.5 Hz), 7.60 (1H, d, J = 2.0 Hz), 7.84 (2H, d, J = 8.2 Hz), 7.89 (2H, d, J = 8.7 Hz), 7.97 (1H, s), 11.39 (1H, s), 12.28 (1H, brs).

IR (Nujol): 1660 cm⁻¹.

m.p.: 132 - 137°C.

<Example 104; Production of 3-(2,4-dichlorobenzyl)-2-methyl-5-((E)-styrenesulfonylcarbamoyl)indole (134)>

According to the method of Example 59, obtained is 3-(2,4-dichlorobenzyl)-2-methyl-5-((E)-styrenesulfonylcarbamoyl)indole (134) (0.242 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.334 g), N,N'-carbonyldiimidazole (0.243 g), (E)-styrenesulfonamide (0.275 g) and diazabicycloundecene (0.228 g).

¹H-NMR (DMSO-d₆, δ ppm): 2.29 (3H, s), 4.08 (2H, s), 6.90 (1H, d, J = 8.4 Hz), 7.25 (1H, dd, J = 8.4 and 2.0 Hz), 7.33 (1H, d, J = 8.6 Hz), 7.45 (3H, m), 7.49 (1H, d, J = 15.4 Hz), 7.61 (3H, m), 7.76 (2H, m), 8.05 (1H, s), 11.38 (1H, s), 11.79 (1H, brs).

IR (Nujol): 1674 cm⁻¹.

m.p.: 190 - 196°C.

<Example 105; Production of 3-(2,4-dichlorobenzyl)-2-methyl-5-(4-vinylbenzene)sulfonylcarbamoyl)indole (135)>

According to the method of Example 59, obtained is 3-(2,4-dichlorobenzyl)-2-methyl-5-(4-vinylbenzene)sulfonylcarbamoyl)indole (135) (0.175 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.334 g), N,N'-carbonyldiimidazole (0.243 g), (4-vinylbenzene)sulfonamide (0.275 g) and diazabicycloundecene (0.228 g).

¹H-NMR (DMSO-d₆, δ ppm): 2.28 (3H, s), 4.07 (2H, s), 5.45 (1H, d, J = 10.9 Hz), 6.01 (1H, d, J = 17.6 Hz), 6.81 (1H, dd, J = 11.0 and 17.6 Hz), 6.90 (1H, d, J = 8.4 Hz), 7.25 (1H, m), 7.31 (1H, d, J = 8.5 Hz), 7.54 (1H, d, J = 8.4 Hz), 7.60 (1H, s), 7.69 (2H, d, J = 8.2 Hz), 7.93 (2H, d, J = 8.2 Hz), 7.97 (1H, s), 11.37 (1H, s), 12.16 (1H, brs).

IR (Nujol): 1674 cm⁻¹.

m.p.: 143 - 146°C.

<Example 106; Production of 5-((4-phenylazobenzene)sulfonylcarbamoyl)-3-(2,4-dichlorobenzyl)-2-methylindole (136)>

5-Carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.334 g) and N,N'-carbonyldiimidazole (0.243 g) are dissolved in N,N-dimethylformamide (5 ml) and stirred at room temperature for 90 minutes. Next, (4-phenylazobenzene)sulfonamide (0.392 g) and diazabicycloundecene (0.228 g) are added thereto, and stirred at 100°C for 7 hours. The reaction mixture is cooled to room temperature, to which are added chloroform (40 ml), water (40 ml) and 1 N HCl (3 ml) for liquid-liquid separation. The organic layer is concentrated under reduced pressure, and the resulting residue is again subjected to liquid-liquid separation with ethyl acetate (40 ml) and water (40 ml) added thereto. Then, the organic layer is dried with anhydrous sodium sulfate, and concentrated under reduced pressure. The resulting residue is recrystallized from chloroform (4 ml) and t-butyl methyl ether (2 ml) to obtain 5-((4-phenylazobenzene)sulfonylcarbamoyl)-3-(2,4-dichlorobenzyl)-2-methylindole (136) (0.150 g).

¹H-NMR (DMSO-d₆, δ ppm): 2.28 (3H, s), 4.07 (2H, s), 6.91 (1H, d, J = 8.4 Hz), 7.25 (1H, dd, J = 2.1 and 8.3 Hz), 7.32 (1H, d, J = 8.6 Hz), 7.56 (1H, dd, J = 1.7 and 8.6 Hz), 7.60 (1H, d, J = 2.3 Hz), 7.62 (3H, m), 7.93 (2H, m), 8.00 (1H, m), 8.06 (2H, d, J = 8.7 Hz), 8.18 (2H, d, J = 8.7 Hz), 11.39 (1H, s), 12.34 (1H, brs).

IR (Nujol): 1677 cm⁻¹.

m.p.: 207 - 223°C.

<Example 107; Production of 3-(2,4-dichlorobenzyl)-2-methyl-5-((4-trifluoromethylbenzene)sulfonylcarbamoyl)indole (137)>

According to the method of Example 106, obtained is 3-(2,4-dichlorobenzyl)-2-methyl-5-((4-trifluoromethylbenzene)sulfonylcarbamoyl)indole (137) (0.360 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.334 g), N,N'-carbonyldiimidazole (0.243 g), (4-trifluoromethylbenzene)sulfonamide (0.338 g) and diazabicycloundecene (0.228 g).

$^1\text{H-NMR}$ (DMSO- d_6 , δ ppm): 2.28 (3H, s), 4.07 (2H, s), 6.90 (1H, d, $J = 8.4$ Hz), 7.25 (1H, dd, $J = 2.2$ and 8.4 Hz), 7.32 (1H, d, $J = 8.6$ Hz), 7.55 (1H, dd, $J = 1.6$ and 8.6 Hz), 7.60 (1H, d, $J = 2.2$ Hz), 7.98 (1H, s), 8.01 (2H, d, $J = 8.3$ Hz), 8.17 (2H, d, $J = 8.3$ Hz), 11.40 (1H, s), 12.41 (1H, brs).

IR (Nujol): 1622 cm^{-1} .

m.p.: $147 - 151^\circ\text{C}$.

<Example 108; Production of 3-(2,4-dichlorobenzyl)-2-methyl-5-((4-methyl-1-pent-1-ene)sulfonylcarbamoyl)indole (138)>

According to the method of Example 106, obtained is 3-(2,4-dichlorobenzyl)-2-methyl-5-((4-methyl-1-pent-1-ene)sulfonylcarbamoyl)indole (138) (0.218 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.400 g), N,N'-carbonyldiimidazole (0.388 g), (4-methyl-1-pent-1-ene)sulfonamide (0.391 g) and diazabicycloundecene (0.364 g).

$^1\text{H-NMR}$ (DMSO- d_6 , δ ppm): 0.86 (6H, d, $J = 6.6$ Hz), 1.71 - 1.80 (1H, m), 2.14 (2H, t, $J = 6.7$ Hz), 2.30 (3H, s), 4.08 (2H, s), 6.74 - 6.85 (2H, m), 6.92 (1H, d, $J = 8.4$ Hz), 7.26 (1H, dd, $J = 2.1$ and 8.4 Hz), 7.34 (1H, d, $J = 8.5$ Hz), 7.59 (1H, d, $J = 1.3$ Hz), 7.61 (1H, d, $J = 1.3$ Hz), 8.00 (1H, d, $J = 1.3$ Hz), 11.38 (1H, s), 11.79 (1H, s).

IR (Nujol): 1682 cm^{-1} .

m.p.: $199.0 - 203.0^\circ\text{C}$.

<Example 109; Production of 3-(2,4-dichlorobenzyl)-5-((3,4-dimethoxybenzene)sulfonylcarbamoyl)-2-methylindole (139)>

According to the method of Example 106, obtained is 3-(2,4-dichlorobenzyl)-5-((3,4-dimethoxybenzene)sulfonylcarbamoyl)-2-methylindole (139) (0.167 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.334 g), N,N'-carbonyldiimidazole (0.243 g), (3,4-dimethoxybenzene)sulfonamide (0.326 g) and diazabicycloundecene (0.228 g).

¹H-NMR (DMSO-d₆, δ ppm): 2.28 (3H, s), 3.79 (3H, s), 3.83 (3H, s), 4.07 (2H, s), 6.90 (1H, d, J = 8.4 Hz), 7.15 (1H, d, J = 8.6 Hz), 7.25 (1H, dd, J = 2.0 and 8.4 Hz), 7.31 (1H, d, J = 8.6 Hz), 7.45 (1H, d, J = 2.1 Hz), 7.54 (1H, dd, J = 1.5 and 8.6 Hz), 7.57 (1H, dd, J = 2.1 and 8.6 Hz), 7.60 (1H, d, J = 2.1 Hz), 7.96 (1H, s), 11.36 (1H, s), 11.97 (1H, s).

IR (Nujol): 1673 cm⁻¹.

m.p.: 148 - 152°C.

<Example 110; Production of 5-((4-t-butylbenzene)sulfonylcarbamoyl)-3-(2,4-dichlorobenzyl)-2-methylindole (140)>

According to the method of Example 106, obtained is 5-((4-t-butylbenzene)sulfonylcarbamoyl)-3-(2,4-dichlorobenzyl)-2-methylindole (140) (0.123 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.334 g), N,N'-carbonyldiimidazole (0.243 g), (4-t-butylbenzene)sulfonamide (0.320 g) and diazabicycloundecene (0.228 g).

¹H-NMR (DMSO-d₆, δ ppm): 1.28 (9H, s), 2.28 (3H, s), 4.07 (2H, s), 6.91 (1H, d, J = 7.9 Hz), 7.25 (1H, d, J = 8.3 Hz), 7.31 (1H, d, J = 8.1 Hz), 7.54 (1H, d, J = 8.2 Hz), 7.61 (3H, m), 7.89 (2H, d, J = 7.3 Hz), 7.98 (1H, s), 11.37 (1H, s), 12.12 (1H, brs).

IR (Nujol): 1688 cm^{-1} .

m.p.: 203 - 207°C.

<Example 111; Production of 3-(2,4-dichlorobenzyl)-2-methyl-5-((3-methylbenzene)sulfonylcarbamoyl)indole (141)>

According to the method of Example 106, obtained is 3-(2,4-dichlorobenzyl)-2-methyl-5-((3-methylbenzene)sulfonylcarbamoyl)indole (141) (0.200 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.250 g), N,N'-carbonyldiimidazole (0.182 g), (3-methylbenzene)sulfonamide (0.192 g) and diazabicycloundecene (0.171 g).

$^1\text{H-NMR}$ (DMSO- d_6 , δ ppm): 2.28 (3H, s), 2.38 (3H, s), 4.07 (2H, s), 6.90 (1H, d, $J = 8.4$ Hz), 7.25 (1H, dd, $J = 2.1$ and 8.3 Hz), 7.30 (1H, d, $J = 8.5$ Hz), 7.46 - 7.50 (2H, m), 7.55 (1H, d, $J = 8.5$ Hz), 7.60 (1H, d, $J = 2.1$ Hz), 7.73 - 7.78 (2H, m), 7.97 (1H, s), 11.36 (1H, s), 12.14 (1H, brs).

IR (Nujol): 1683 cm^{-1} .

m.p.: 105 - 115°C (foamy solid).

<Example 112; Production of 3-(2,4-dichlorobenzyl)-2-methyl-5-(2-octanesulfonylcarbamoyl)indole (142)>

According to the method of Example 106, obtained is 3-(2,4-dichlorobenzyl)-2-methyl-5-(2-octanesulfonylcarbamoyl)indole (142) (0.032 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.107 g), N,N'-carbonyldiimidazole (0.063 g), 2-octanesulfonamide (0.075 g) and diazabicycloundecene (0.059 g).

$^1\text{H-NMR}$ (DMSO- d_6 , δ ppm): 0.85 (3H, t, $J = 7.7$ Hz), 1.2 - 1.5 (8H, m), 1.43 (3H, d, $J = 6.9$ Hz), 1.64 (1H, m), 2.05 (1H, m), 2.38 (3H, s), 3.89 (1H, m), 4.11 (2H, s), 6.80 (1H, d, $J = 8.3$ Hz), 7.04 (1H, dd, $J = 2.0$ and 8.3 Hz), 7.36 (1H, d, $J = 8.5$ Hz), 7.42 (1H, d, $J = 2.0$ Hz), 7.60 (1H, dd, $J = 1.6$ and 8.5 Hz), 7.84 (1H, brs), 8.21 (2H, brs).

IR (Nujol): 1678 cm^{-1} .

m.p.: 181 - 182°C.

<Example 113; Production of 3-(2,4-dichlorobenzyl)-2-methyl-5-((4-phenylbenzene)sulfonylcarbamoyl)indole (143)>

According to the method of Example 106, obtained is 3-(2,4-dichlorobenzyl)-2-methyl-5-((4-phenylbenzene)sulfonylcarbamoyl)indole (143) (0.109 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.334 g), N,N'-carbonyldiimidazole (0.243 g), (4-phenylbenzene)sulfonamide (0.350 g) and diazabicycloundecene (0.228 g).

^1H -NMR (DMSO- d_6 , δ ppm): 2.28 (3H, s), 4.07 (2H, s), 6.91 (1H, d, $J = 8.4$ Hz), 7.25 (1H, dd, $J = 2.0$ and 8.3 Hz), 7.32 (1H, d, $J = 8.6$ Hz), 7.43 (1H, t, $J = 7.3$ Hz), 7.50 (2H, t, $J = 7.5$ Hz), 7.55 (1H, dd, $J = 1.3$ and 8.6 Hz), 7.60 (1H, d, $J = 1.9$ Hz), 7.73 (2H, d, $J = 8.6$ Hz), 7.90 (2H, d, $J = 8.5$ Hz), 7.99 (1H, s), 8.04 (2H, d, $J = 8.5$ Hz), 11.38 (1H, s), 12.22 (1H, brs).

IR (Nujol): 1678 cm^{-1} .

m.p.: 206 - 207°C.

<Example 114; Production of 3-(2-chloro-4-phenylbenzyl)-5-methoxycarbonyl-2-methylindole (144)>

A mixture of 5-methoxycarbonyl-2-methylindole (0.84 g), 2-chloro-4-phenylbenzyl bromide (2.0 g), L-tartaric acid (1.07 g), sodium hydroxide (0.37 g), sodium iodide (0.53 g), 1,4-dioxane (15 ml) and water (8 ml) is stirred at 90°C for 4 hours. The reaction mixture is extracted with water and ethyl acetate added thereto, and the organic layer is washed with water, dried and concentrated. The resulting residue is purified through silica gel column chromatography (eluent: ethyl acetate/hexane = 1/4 to 1/2), and crystallized from a mixed solution of hexane and ethyl acetate to obtain 3-(2-chloro-4-phenylbenzyl)-5-methoxycarbonyl-2-methylindole (144) (0.965).

¹H-NMR (DMSO-d₆, δ ppm): 2.38 (3H, s), 3.77 (3H, s), 4.16 (2H, s), 7.04 (1H, d, J = 8.1 Hz), 7.36 (2H, t, J = 5.9 Hz), 7.44 (2H, t, J = 7.2 Hz), 7.48 (1H, d, J = 8.0 Hz), 7.65 (3H, m), 7.74 (1H, s), 7.96 (1H, s), 11.36 (1H, s).

<Example 115; Production of 5-carboxy-3-(2-chloro-4-phenylbenzyl)-2-methylindole (145)>

To 3-(2-chloro-4-phenylbenzyl)-5-methoxycarbonyl-2-methylindole (0.95 g), added are ethanol (5 ml), water (5 ml) and aqueous sodium hydroxide (0.29 g), and refluxed for 3 hours. The reaction mixture is cooled, and neutralized with HCl. This is extracted with ethyl acetate and water, and the organic layer is dried and concentrated. The resulting residue is crystallized from a small amount of a mixed solution of chloroform and ether. The crystal is taken out through filtration and dried to obtain 5-carboxy-3-(2-chloro-4-phenylbenzyl)-2-methylindole (145) (0.701 g).

¹H-NMR (DMSO-d₆, δ ppm): 2.37 (3H, s), 4.13 (2H, s), 7.04 (1H, d, J = 8.1 Hz), 7.28 (1H, d, J = 8.5 Hz), 7.36 (1H, d, J = 6.8 Hz), 7.44 (2H, d, J = 7.6 Hz), 7.47 (1H, d, J = 8.0 Hz), 7.64 (3H, m), 7.73 (1H, d, J = 1.8 Hz), 7.92 (1H, s), 11.24 (1H, s).

<Example 116; Production of 3-((2-chloro-4-phenyl)benzyl)-2-methyl-5-((1-pent-1-ene)sulfonylcarbamoyl)indole (146)>

According to the method of Example 106, obtained is 3-((2-chloro-4-phenyl)benzyl)-2-methyl-5-((1-pent-1-ene)sulfonylcarbamoyl)indole (146) (0.135 g) from 5-carboxy-3-((2-chloro-4-phenyl)benzyl)-2-methylindole (0.301 g), N,N'-carbonyldiimidazole (0.195 g), (1-pent-1-ene)sulfonamide (0.179 g) and diazabicycloundecene (0.183 g).

¹H-NMR (DMSO-d₆, δ ppm): 0.84 (3H, t, J = 7.4 Hz), 1.42 (2H, m), 2.21 (2H, m), 2.33 (3H, s), 4.14 (2H, s), 6.75 (1H, d, J = 15.2 Hz), 6.83 (1H, dt, J = 6.4 and

15.1 Hz), 6.98 (1H, d, $J = 8.0$ Hz), 7.35 (2H, m), 7.42 - 7.47 (3H, m), 7.60 - 7.64 (3H, m), 7.74 (1H, d, $J = 1.7$ Hz), 8.07 (1H, s), 11.38 (1H, s), 11.81 (1H, s).

IR (Nujol): 1654 cm^{-1} .

m.p.: $166 - 167^{\circ}\text{C}$.

<Example 117; Production of 3-(2-chloro-4-phenylbenzyl)-2-methyl-5-(1-pentanesulfonylcarbamoyl)indole (147)>

According to the method of Example 106, obtained is 3-(2-chloro-4-phenylbenzyl)-2-methyl-5-(1-pentanesulfonylcarbamoyl)indole (147) (0.122 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.200 g), N,N'-carbonyldiimidazole (0.173 g), 1-pentanesulfonamide (0.165 g) and diazabicycloundecene (0.162 g).

$^1\text{H-NMR}$ (DMSO- d_6 , δ ppm): 0.79 (3H, t, $J = 7.3$ Hz), 1.21 - 1.41 (4H, m), 1.65 - 1.72 (2H, m), 2.34 (3H, s), 3.48 (2H, t, $J = 7.7$ Hz), 4.16 (2H, s), 7.01 (1H, d, $J = 8.1$ Hz), 7.37 (2H, d, $J = 8.2$ Hz), 7.42 - 7.49 (3H, m), 7.64 (3H, m), 7.75 (1H, d, $J = 1.8$ Hz), 8.10 (1H, s), 11.40 (1H, s), 11.71 (1H, s).

IR (Nujol): 1684 cm^{-1} .

m.p.: $155.5 - 156.5^{\circ}\text{C}$.

<Example 118; Production of 5-(benzenesulfonylcarbamoyl)-3-((2-chloro-4-phenyl)benzyl)-2-methylindole (148)>

According to the method of Example 106, obtained is 5-(benzenesulfonylcarbamoyl)-3-((2-chloro-4-phenyl)benzyl)-2-methylindole (148) (0.057 g) from 5-carboxy-3-((2-chloro-4-phenyl)benzyl)-2-methylindole (0.225 g), N,N'-carbonyldiimidazole (0.221 g), benzenesulfonamide (0.214 g) and diazabicycloundecene (0.207 g).

$^1\text{H-NMR}$ (DMSO- d_6 , δ ppm): 2.32 (3H, s), 4.13 (2H, s), 6.97 (1H, d, $J = 8.0$ Hz), 7.31 - 7.37 (2H, m), 7.42 - 7.47 (3H, m), 7.54 - 7.67 (6H, m), 7.73 (1H, s),

7.96 (2H, d, $J = 8.0$ Hz), 8.03 (1H, s), 11.38 (1H, s), 12.20 (1H, brs).

IR (Nujol): 1661 cm^{-1} .

m.p.: $134 - 137^{\circ}\text{C}$.

<Example 119; Production of 3-(2,4-dichlorobenzyl)-5-((4-ethylbenzene)sulfonylcarbamoyl)-2-methylindole (149)>

According to the method of Example 106, obtained is 3-(2,4-dichlorobenzyl)-5-((4-ethylbenzene)sulfonylcarbamoyl)-2-methylindole (149) (0.108 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.334 g), N,N' -carbonyldiimidazole (0.243 g), (4-ethylbenzene)sulfonamide (0.278 g) and diazabicycloundecene (0.228 g).

$^1\text{H-NMR}$ (DMSO- d_6 , δ ppm): 1.18 (3H, t, $J = 7.5$ Hz), 2.28 (3H, s), 2.67 (2H, q, $J = 7.3$ Hz), 4.07 (2H, s), 6.91 (1H, d, $J = 8.3$ Hz), 7.25 (1H, d, $J = 7.5$ Hz), 7.31 (1H, d, $J = 8.4$ Hz), 7.44 (2H, d, $J = 7.9$ Hz), 7.54 (1H, d, $J = 8.6$ Hz), 7.60 (1H, s), 7.87 (2H, d, $J = 8.0$ Hz), 7.97 (1H, s), 11.37 (1H, s), 12.10 (1H, brs).

IR (Nujol): 1682 cm^{-1} .

m.p.: $197 - 198^{\circ}\text{C}$.

<Example 120; Production of 5-((4-n-butylbenzene)sulfonylcarbamoyl)-3-(2,4-dichlorobenzyl)-2-methylindole (150)>

According to the method of Example 106, obtained is 5-((4-n-butylbenzene)sulfonylcarbamoyl)-3-(2,4-dichlorobenzyl)-2-methylindole (150) (0.230 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.334 g), N,N' -carbonyldiimidazole (0.243 g), (4-n-butylbenzene)sulfonamide (0.320 g) and diazabicycloundecene (0.228 g).

$^1\text{H-NMR}$ (DMSO- d_6 , δ ppm): 0.87 (3H, t, $J = 7.3$ Hz), 1.28 (2H, m), 1.55 (2H, m), 2.28 (3H, s), 2.64 (2H, t, $J = 7.6$ Hz), 4.07 (2H, s), 6.90 (1H, d, $J = 8.4$ Hz), 7.25 (1H, dd, $J = 2.0$ and 8.3 Hz), 7.31 (1H, d, $J = 8.6$ Hz), 7.42 (1H, d, $J = 8.2$

Hz), 7.54 (1H, dd, $J = 1.5$ and 8.6 Hz), 7.60 (1H, d, $J = 2.0$ Hz), 7.87 (2H, d, $J = 8.3$ Hz), 7.97 (1H, s), 11.37 (1H, s), 12.09 (1H, s).

IR (Nujol): 1659 cm^{-1} .

m.p.: $154 - 155^{\circ}\text{C}$.

<Example 121; Production of 5-((4-n-butoxybenzene)sulfonylcarbamoyl)-3-(2,4-dichlorobenzyl)-2-methylindole (151)>

According to the method of Example 106, obtained is 5-((4-n-butoxybenzene)sulfonylcarbamoyl)-3-(2,4-dichlorobenzyl)-2-methylindole (151) (0.140 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.334 g), N,N'-carbonyldiimidazole (0.243 g), (4-n-butoxybenzene)sulfonamide (0.344 g) and diazabicycloundecene (0.228 g).

^1H -NMR (DMSO- d_6 , δ ppm): 0.91 (3H, t, $J = 7.1$ Hz), 1.42 (2H, m), 1.69 (2H, m), 2.28 (3H, s), 4.04 (2H, t, $J = 6.5$ Hz), 4.06 (2H, s), 6.90 (1H, d, $J = 8.3$ Hz), 7.10 (2H, d, $J = 8.0$ Hz), 7.24 (1H, d, $J = 8.0$ Hz), 7.30 (1H, d, $J = 8.5$ Hz), 7.54 (1H, d, $J = 8.5$ Hz), 7.60 (1H, s), 7.88 (2H, d, $J = 7.9$ Hz), 7.96 (1H, s), 11.36 (1H, s), 12.04 (1H, brs).

IR (Nujol): 1654 cm^{-1} .

m.p.: $185 - 189^{\circ}\text{C}$.

<Example 122; Production of 3-(2,4-dichlorobenzyl)-5-methoxycarbonyl-2-methylthioindole (152)>

To a methylene chloride (10 ml) solution of trifluoroacetic acid (1.238 g) and triethylsilane (2.52 g), added is a methylene chloride (10 ml) solution of 5-methoxycarbonyl-2-methylthioindole (0.800 g), which had been prepared according to the method described in "Tetrahedron, 42 (16), 4511 (1986)", and 2,4-dichlorobenzaldehyde (0.760 g), in an ice-water bath. This is stirred for 3 hours in the

ice-water bath. The crystal formed is taken out through filtration and dried to obtain 3-(2,4-dichlorobenzyl)-5-methoxycarbonyl-2-methylthioindole (152) (0.727 g).

$^1\text{H-NMR}$ (CDCl_3 , δ ppm): 2.33 (3H, s), 3.89 (3H, s), 4.29 (2H, s), 6.79 (1H, d, J = 8.3 Hz), 7.02 (1H, d, J = 8.3 Hz), 7.35 (1H, d, J = 8.6 Hz), 7.42 (1H, d, J = 1.9 Hz), 7.92 (1H, dd, J = 1.3 and 8.7 Hz), 8.10 (1H, s), 8.32 (1H, brs).

<Example 123; Production of 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylthioindole (153)>

According to the method of Example 115, obtained is 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylthioindole (153) (0.795 g) from 3-(2,4-dichlorobenzyl)-5-methoxycarbonyl-2-methylthioindole (0.950 g).

$^1\text{H-NMR}$ (DMSO-d_6 , δ ppm): 2.44 (3H, s), 4.20 (2H, s), 6.94 (1H, d, J = 8.3 Hz), 7.28 (1H, dd, J = 2.2 and 8.4 Hz), 7.38 (1H, d, J = 8.6 Hz), 7.62 (1H, d, J = 2.2 Hz), 7.72 (1H, d, J = 8.7 Hz), 7.92 (1H, s), 11.79 (1H, s), 12.44 (1H, brs).

<Example 124; Production of 3-(2,4-dichlorobenzyl)-2-methylthio-5-((1-pent-1-ene)sulfonylcarbamoyl)indole (154)>

According to the method of Example 106, obtained is 3-(2,4-dichlorobenzyl)-2-methylthio-5-((1-pent-1-ene)sulfonylcarbamoyl)indole (154) (0.068 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylthioindole (0.200 g), N,N' -carbonyldiimidazole (0.177 g), (1-pent-1-ene)sulfonamide (0.163 g) and diazabicycloundecene (0.166 g).

$^1\text{H-NMR}$ (DMSO-d_6 , δ ppm): 0.86 (3H, t, J = 7.4 Hz), 1.40 - 1.49 (2H, m), 2.23 (2H, q, J = 7.2 Hz), 2.41 (3H, s), 4.18 (2H, s), 6.76 (1H, d, J = 6.0 Hz), 6.82 - 6.89 (2H, m), 7.26 (1H, dd, J = 1.9 and 8.4 Hz), 7.40 (1H, d, J = 8.7 Hz), 8.63 (1H, d, J = 2.0 Hz), 7.70 (1H, d, J = 8.8 Hz), 8.07 (1H, s), 11.86 (1H, brs), 11.88 (1H, brs).

IR (Nujol): 1682 cm^{-1} .

m.p.: 140.5 - 142.0°C.

<Example 125; Production of 5-(benzenesulfonylcarbamoyl)-3-(2,4-dichlorobenzyl)-2-methylthioindole (155)>

According to the method of Example 106, obtained is 5-(benzenesulfonylcarbamoyl)-3-(2,4-dichlorobenzyl)-2-methylthioindole (155) (0.100 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylthioindole (0.160 g), N,N'-carbonyldiimidazole (0.092 g), benzenesulfonamide (0.089 g) and diazabicycloundecene (0.086 g).

¹H-NMR (DMSO-d₆, δ ppm): 2.41 (3H, s), 4.17 (2H, s), 6.82 (1H, d, J = 8.4 Hz), 7.26 (1H, dd, J = 2.1 and 8.3 Hz), 7.38 (1H, d, J = 8.7 Hz), 7.60 - 7.66 (4H, m), 7.70 (1H, t, J = 7.3 Hz), 7.98 (2H, d, J = 7.5 Hz), 8.04 (1H, s), 11.88 (1H, brs), 12.26 (1H, brs).

IR (Nujol): 1698 cm⁻¹.

m.p.: 223.0 - 226.0°C.

<Example 126; Production of 3-(2,4-dichlorobenzyl)-2-methylthio-5-(1-pentanesulfonylcarbamoyl)indole (156)>

According to the method of Example 106, obtained is 3-(2,4-dichlorobenzyl)-2-methylthio-5-(1-pentanesulfonylcarbamoyl)indole (156) (0.218 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylthioindole (0.200 g), N,N'-carbonyldiimidazole (0.177 g), 1-pentanesulfonamide (0.165 g) and diazabicycloundecene (0.166 g).

¹H-NMR (DMSO-d₆, δ ppm): 0.81 (3H, t, J = 7.3 Hz), 1.22 - 1.31 (2H, m), 1.31 - 1.39 (2H, m), 1.63 - 1.70 (2H, m), 2.42 (3H, d, J = 0.7 Hz), 3.49 (2H, t, J = 7.5 Hz), 4.19 (2H, s), 6.85 (1H, d, J = 8.4 Hz), 7.26 (1H, d, J = 8.4 Hz), 7.42 (1H, d, J = 8.7 Hz), 8.63 (1H, d, J = 1.9 Hz), 7.72 (1H, d, J = 8.6 Hz), 8.09 (1H, s), 11.76 (1H, s), 11.89 (1H, s).

IR (Nujol): 1682 cm^{-1} .

m.p.: 176.5 - 180.0°C.

<Example 127: Production of 3-(2,4-dichlorobenzyl)-2-methyl-5-((1-penta-1,3-diene)sulfonylcarbamoyl)indole (157)>

According to the method of Example 106, obtained is 3-(2,4-dichlorobenzyl)-2-methyl-5-((1-penta-1,3-diene)sulfonylcarbamoyl)indole (157) (0.174 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylindole (0.300 g), N,N'-carbonyldiimidazole (0.291 g), (1-penta-1,3-diene)sulfonamide (0.264 g) and diazabicycloundecene (0.273 g).

^1H -NMR (DMSO- d_6 , δ ppm): 1.83 (3H, d, $J = 6.4$ Hz), 2.30 (3H, s), 4.09 (2H, s), 6.25 - 6.42 (2H, m), 6.77 (1H, d, $J = 14.8$ Hz), 6.92 (1H, d, $J = 8.4$ Hz), 7.18 (1H, dd, $J = 9.5$ and 14.8 Hz), 7.26 (1H, dd, $J = 2.2$ and 8.4 Hz), 7.34 (1H, d, $J = 8.5$ Hz), 7.61 (2H, m), 8.02 (1H, s), 11.38 (1H, brs), 11.82 (1H, brs).

IR (Nujol): 1682 cm^{-1} .

m.p.: 174.0 - 177.0°C.

<Example 128; Production of 5-((2-cyclopropylethylene)sulfonylcarbamoyl)-3-(2,4-dichlorobenzyl)-2-methylindole (158)>

According to the method of Example 106, obtained is 5-((2-cyclopropylethylene)sulfonylcarbamoyl)-3-(2,4-dichlorobenzyl)-2-methylindole (158) (0.180 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylthioindole (0.300 g), N,N'-carbonyldiimidazole (0.291 g), (2-cyclopropylethylene)sulfonylcarbamoyl (0.264 g) and diazabicycloundecene (0.273 g).

^1H -NMR (DMSO- d_6 , δ ppm): 0.70 - 0.74 (2H, m), 0.94 - 0.98 (2H, m), 1.68 - 1.77 (1H, m), 2.30 (3H, s), 4.09 (2H, s), 6.34 (1H, dd, $J = 10.4$ and 14.8 Hz), 6.84 (1H, d, $J = 14.8$ Hz), 6.92 (1H, d, $J = 8.4$ Hz), 7.27 (1H, dd, $J = 2.2$ and 8.4

Hz), 7.34 (1H, d, $J = 8.6$ Hz), 7.62 (2H, m), 8.02 (1H, s), 11.38 (1H, brs), 11.73 (1H, brs).

IR (Nujol): 1684 cm^{-1} .

m.p.: $172.5 - 174.0^{\circ}\text{C}$.

<Example 129; Production of 3-(2,4-dichlorobenzyl)-2-methyl-5-((4-methyl-(E)-styrene)sulfonylcarbamoyl)indole (159)>

According to the method of Example 106, obtained is 3-(2,4-dichlorobenzyl)-2-methyl-5-((4-methyl-(E)-styrene)sulfonylcarbamoyl)indole (159) (0.122 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylthioindole (0.200 g), N,N'-carbonyldiimidazole (0.126 g), (4-methyl-(E)-styrene)sulfonamide (0.153 g) and diazabicycloundecene (0.118 g).

$^1\text{H-NMR}$ (DMSO- d_6 , δ ppm): 2.29 (3H, s), 2.34 (3H, s), 4.08 (2H, s), 6.91 (1H, d, $J = 8.4$ Hz), 7.26 (3H, m), 7.34 (1H, d, $J = 8.5$ Hz), 7.42 (1H, d, $J = 15.4$ Hz), 7.56 - 7.67 (5H, m), 8.05 (1H, s), 11.38 (1H, brs), 11.95 (1H, brs).

IR (Nujol): 1678 cm^{-1} .

m.p.: $253.5 - 255.0^{\circ}\text{C}$.

<Example 130; Production of 3-(2,4-dichlorobenzyl)-5-((4-methoxy-(E)-styrene)sulfonylcarbamoyl)-2-methylindole (160)>

According to the method of Example 106, obtained is 3-(2,4-dichlorobenzyl)-5-((4-methoxy-(E)-styrene)sulfonylcarbamoyl)-2-methylindole (160) (0.067 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methylthioindole (0.200 g), N,N'-carbonyldiimidazole (0.126 g), (4-methoxy-(E)-styrene)sulfonamide (0.165 g) and diazabicycloundecene (0.118 g).

$^1\text{H-NMR}$ (DMSO- d_6 , δ ppm): 2.29 (3H, s), 3.80 (3H, s), 4.08 (2H, s), 6.91 (1H, d, $J = 8.5$ Hz), 6.99 (2H, d, $J = 8.8$ Hz), 7.25 (1H, dd, $J = 8.4$ and 2.1 Hz), 7.32

(1H, d, J = 15.4 Hz), 7.33 (1H, d, J = 8.4 Hz), 7.55 (1H, d, J = 15.4 Hz), 7.60 - 7.64 (2H, m), 7.71 (2H, d, J = 8.7 Hz), 8.05 (1H, s), 11.37 (1H, s), 11.89 (1H, brs).

IR (Nujol): 1674 cm^{-1} .

m.p.: 227.5 - 231.0°C.

<Example 131; Production of 3-(2,4-dichlorobenzyl)-5-methoxycarbonyl-2-(methoxymethyl)indole (161)>

Trifluoroacetic acid (1.14 g) and triethylsilane (2.33 g) are dissolved in dichloromethane (10 ml), and the resulting solution is cooled at 0°C, to which is dropwise added a solution of 5-methoxycarbonyl-2-(methoxymethyl)indole (1.10 g) and 2,4-dichlorobenzaldehyde (0.96 g) dissolved in dichloromethane (20 ml). The reaction mixture is stirred at 0°C for 3 hours, and thereafter an aqueous solution of 10 % sodium hydroxide (5 ml), chloroform (40 ml) and a saturated saline solution (40 ml) are added thereto in that order for liquid-liquid separation. The organic layer is dried with anhydrous sodium sulfate, and concentrated under reduced pressure. The resulting residue is purified through silica gel column chromatography (ethyl acetate/hexane = 3/7) to obtain 3-(2,4-dichlorobenzyl)-5-methoxycarbonyl-2-(methoxymethyl)indole (161) (0.61 g).

^1H -NMR (DMSO- d_6 , δ ppm): 3.37 (3H, s), 3.90 (3H, s), 4.16 (2H, s), 4.54 (2H, s), 6.86 (1H, d, J = 8.4 Hz), 7.04 (1H, dd, J = 2.2 and 8.4 Hz), 7.33 (1H, d, J = 9.6 Hz), 7.41 (1H, d, J = 2.2 Hz), 7.90 (1H, dd, J = 1.5 and 8.5 Hz), 8.15 (1H, s), 8.53 (1H, brs).

<Example 132; Production of 5-carboxy-3-(2,4-dichlorobenzyl)-2-(methoxymethyl)indole (162)>

According to the method of Example 115, obtained is 5-carboxy-3-(2,4-

dichlorobenzyl)-2-(methoxymethyl)indole (162) (0.56 g) from 3-(2,4-dichlorobenzyl)-5-methoxycarbonyl-2-(methoxymethyl)indole (0.61 g).

<Example 133; Production of 3-(2,4-dichlorobenzyl)-2-methoxymethyl-5-(1-pentanesulfonylcarbamoyl)indole (163)>

According to the method of Example 106, obtained is 3-(2,4-dichlorobenzyl)-2-methoxymethyl-5-(1-pentanesulfonylcarbamoyl)indole (163) (0.115 g) from 5-carboxy-3-(2,4-dichlorobenzyl)-2-methoxymethylindole (0.364 g), N,N'-carbonyldiimidazole (0.243 g), 1-pentanesulfonamide (0.227 g) and diazabicycloundecene (0.228 g).

¹H-NMR (DMSO-d₆, δ ppm): 0.80 (3H, t, J = 7.3 Hz), 1.25 (2H, m), 1.34 (2H, m), 1.66 (2H, m), 3.32 (3H, s), 3.48 (2H, t, J = 7.7 Hz), 4.17 (2H, s), 4.51 (2H, s), 6.92 (1H, d, J = 8.4 Hz), 7.25 (1H, dd, J = 2.1 and 8.3 Hz), 7.41 (1H, d, J = 8.6 Hz), 7.62 (1H, d, J = 2.1 Hz), 7.69 (1H, dd, J = 1.5 and 8.6 Hz), 8.10 (1H, s), 11.64 (1H, s), 11.73 (1H, s).

IR (Nujol): 1685 cm⁻¹.

m.p.: 189 - 192°C.

<Example 134; Production of 3-((1-bromonaphthalen-2-yl)methyl)-2-methyl-5-((E)-styrene)sulfonylcarbamoyl)indole (164)>

According to the method of Example 106, obtained is 3-((1-bromonaphthalen-2-yl)methyl)-2-methyl-5-((E)-styrene)sulfonylcarbamoyl)indole (164) (0.080 g) from 3-((1-bromonaphthalen-2-yl)methyl)-5-carboxy-2-methylindole (0.237 g), N,N'-carbonyldiimidazole (0.162 g), (E)-styrenesulfonamide (0.183 g) and diazabicycloundecene (0.152 g).

¹H-NMR (DMSO-d₆, δ ppm): 2.31 (3H, s), 4.37 (2H, s), 7.10 (1H, d, J = 8.5 Hz), 7.34 (1H, d, J = 8.6 Hz), 7.43 (3H, m), 7.47 (1H, d, J = 15.4 Hz), 7.55 (1H, t, J = 7.7 Hz), 7.60 (1H, d, J = 15.9 Hz), 7.62 (1H, m), 7.67 (1H, m), 7.74 (2H,

m), 7.78 (1H, d, J = 8.5 Hz), 7.88 (1H, d, J = 8.0 Hz), 8.12 (1H, s), 8.26 (1H, d, J = 8.6 Hz), 11.38 (1H, s), 11.97 (1H, s).

IR (Nujol): 1674 cm^{-1} .

m.p.: 235 - 238°C.

<Example 135; Production of 3-((1-bromonaphthalen-2-yl)methyl)-2-methyl-5-(4-vinylbenzene)sulfonylcarbamoylindole (165)>

According to the method of Example 106, obtained is 3-((1-bromonaphthalen-2-yl)methyl)-2-methyl-5-(4-vinylbenzene)sulfonylcarbamoylindole (165) (0.130 g) from 3-((1-bromonaphthalen-2-yl)methyl)-5-carboxy-2-methylindole (0.237 g), N,N'-carbonyldiimidazole (0.162 g), (4-vinylbenzene)sulfonamide (0.183 g) and diazabicycloundecene (0.152 g).

^1H -NMR (DMSO- d_6 , δ ppm): 2.30 (3H, s), 4.36 (2H, s), 5.44 (1H, d, J = 11.0 Hz), 5.99 (1H, d, J = 17.6 Hz), 6.79 (1H, dd, J = 11.1 and 17.7 Hz), 7.10 (1H, d, J = 8.4 Hz), 7.32 (1H, d, J = 8.6 Hz), 7.55 (2H, m), 7.67 (3H, m), 7.78 (1H, d, J = 8.5 Hz), 7.90 (3H, m), 8.04 (1H, s), 8.26 (1H, d, J = 8.4 Hz), 11.38 (1H, s), 12.16 (1H, s).

IR (Nujol): 1674 cm^{-1} .

m.p.: 222 - 224°C.

<Example 136; Production of 3-((1-bromonaphthalen-2-yl)methyl)-2-methyl-5-(p-toluenesulfonylcarbamoylindole (166)>

According to the method of Example 106, obtained is 3-((1-bromonaphthalen-2-yl)methyl)-2-methyl-5-(p-toluenesulfonylcarbamoylindole (166) (0.145 g) from 3-((1-bromonaphthalen-2-yl)methyl)-5-carboxy-2-methylindole (0.237 g), N,N'-carbonyldiimidazole (0.162 g), p-toluenesulfonamide (0.171 g) and diazabicycloundecene (0.152 g).

¹H-NMR (DMSO-d₆, δ ppm): 2.30 (3H, s), 2.35 (3H, s), 4.36 (2H, s), 7.09 (1H, d, J = 8.5 Hz), 7.31 (1H, d, J = 8.6 Hz), 7.37 (2H, d, J = 8.1 Hz), 7.54 (2H, m), 7.67 (1H, t, J = 7.9 Hz), 7.78 (1H, d, J = 8.5 Hz), 7.83 (2H, d, J = 8.2 Hz), 7.89 (1H, d, J = 8.2 Hz), 8.03 (1H, s), 8.26 (1H, d, J = 8.6 Hz), 11.37 (1H, s), 12.09 (1H, brs).

IR (Nujol): 1674 cm⁻¹.

m.p.: 260 - 262°C.

<Example 137; Production of 5-(benzenesulfonylcarbamoyl)-3-((1-bromonaphthalen-2-yl)methyl)-2-methylindole (167)>

According to the method of Example 106, obtained is 5-(benzenesulfonylcarbamoyl)-3-((1-bromonaphthalen-2-yl)methyl)-2-methylindole (167) (0.120 g) from 3-((1-bromonaphthalen-2-yl)methyl)-5-carboxy-2-methylindole (0.237 g), N,N'-carbonyldiimidazole (0.162 g), benzenesulfonamide (0.157 g) and diazabicycloundecene (0.152 g).

¹H-NMR (DMSO-d₆, δ ppm): 2.30 (3H, s), 4.36 (2H, s), 7.09 (1H, d, J = 8.5 Hz), 7.32 (1H, d, J = 8.5 Hz), 7.53 - 7.60 (4H, m), 7.67 (2H, m), 7.78 (1H, d, J = 8.5 Hz), 7.89 (1H, d, J = 8.2 Hz), 7.95 (2H, m), 8.05 (1H, s), 8.26 (1H, d, J = 8.5 Hz), 11.38 (1H, s), 12.18 (1H, brs).

IR (Nujol): 1674 cm⁻¹.

m.p.: 159 - 161°C.

<Example 138; Production of 3-((2-chloro-4-phenyl)benzyl)-2-methyl-5-((E)-styrenesulfonylcarbamoyl)indole (168)>

According to the method of Example 106, obtained is 3-((2-chloro-4-phenyl)benzyl)-2-methyl-5-((E)-styrenesulfonylcarbamoyl)indole (168) (0.107 g) from 5-carboxy-3-((2-chloro-4-phenyl)benzyl)-2-methylindole (0.226 g), N,N'-

carbonyldiimidazole (0.162 g), (E)-styrenesulfonamide (0.183 g) and diazabicycloundecene (0.152 g).

$^1\text{H-NMR}$ (DMSO- d_6 , δ ppm): 2.32 (3H, s), 4.14 (2H, s), 6.98 (1H, d, $J = 8.1$ Hz), 7.35 (2H, m), 7.43 (6H, m), 7.48 (1H, d, $J = 15.6$ Hz), 7.63 (4H, m), 7.74 (3H, m), 8.11 (1H, s), 11.38 (1H, s), 11.99 (1H, brs).

IR (Nujol): 1670 cm^{-1} .

m.p.: $232 - 235^\circ\text{C}$.

<Example 139; Production of 3-((2-chloro-4-phenyl)benzyl)-2-methyl-5-((4-vinylbenzene)sulfonylcarbamoyl)indole (169)>

According to the method of Example 106, obtained is 3-((2-chloro-4-phenyl)benzyl)-2-methyl-5-((4-vinylbenzene)sulfonylcarbamoyl)indole (169) (0.078 g) from 5-carboxy-3-((2-chloro-4-phenyl)benzyl)-2-methylindole (0.226 g), N,N'-carbonyldiimidazole (0.162 g), (4-vinylbenzene)sulfonamide (0.183 g) and diazabicycloundecene (0.152 g).

$^1\text{H-NMR}$ (DMSO- d_6 , δ ppm): 2.32 (3H, s), 4.13 (2H, s), 5.44 (1H, d, $J = 11.0$ Hz), 5.98 (1H, d, $J = 17.6$ Hz), 6.79 (1H, dd, $J = 17.5$ and 10.9 Hz), 6.98 (1H, d, $J = 8.1$ Hz), 7.32 (1H, d, $J = 8.6$ Hz), 7.36 (1H, d, $J = 7.4$ Hz), 7.42 - 7.47 (3H, m), 7.54 (1H, dd, $J = 8.4$ and 2.0 Hz), 7.63 (2H, m), 7.67 (2H, d, $J = 8.5$ Hz), 7.73 (1H, d, $J = 1.8$ Hz), 7.91 (2H, d, $J = 8.5$ Hz), 8.03 (1H, s), 11.37 (1H, s), 12.21 (1H, brs).

IR (Nujol): 1670 cm^{-1} .

m.p.: $204 - 210^\circ\text{C}$.

<Example 140; Production of 3-((1-bromonaphthalen-2-yl)methyl)-2-methyl-5-((1-pent-2-ene)sulfonylcarbamoyl)indole (170)>

According to the method of Example 106, obtained is 3-((1-bromonaphthalen-2-

yl)methyl)-2-methyl-5-((1-pent-2-ene)sulfonylcarbamoyl)indole (170) (0.058 g) from 3-((1-bromonaphthalen-2-yl)methyl-5-carboxy-2-methylindole (0.237 g), N,N'-carbonyldiimidazole (0.162 g), (1-pent-1-ene)sulfonamide (0.149 g) and diazabicycloundecene (0.152 g).

¹H-NMR (DMSO-d₆, δ ppm): 0.83 (3H, brs), 1.41 (2H, m), 2.19 (2H, m), 2.32 (3H, s), 4.37 (2H, s), 6.73 (1H, d, J = 14.9 Hz), 6.80 (1H, m), 7.10 (1H, d, J = 8.1 Hz), 7.34 (1H, d, J = 8.3 Hz), 7.55 (1H, m), 7.60 (1H, d, J = 8.2 Hz), 7.67 (1H, m), 7.78 (1H, d, J = 8.2 Hz), 7.89 (1H, d, J = 7.8 Hz), 8.07 (1H, s), 8.27 (1H, d, J = 8.1 Hz), 11.38 (1H, s), 11.79 (1H, brs).

IR (Nujol): 1670 cm⁻¹.

m.p.: 104 - 113°C.

<Example 141; Production of 3-((2-chloro-4-phenyl)benzyl)-2-methyl-5-(p-toluenesulfonylcarbamoyl)indole (171)>

According to the method of Example 106, obtained is 3-((2-chloro-4-phenyl)benzyl)-2-methyl-5-(p-toluenesulfonylcarbamoyl)indole (171) (0.063 g) from 5-carboxy-3-((2-chloro-4-phenyl)benzyl)-2-methylindole (0.226 g), N,N'-carbonyldiimidazole (0.162 g), p-toluenesulfonamide (0.171 g) and diazabicycloundecene (0.152 g).

¹H-NMR (DMSO-d₆, δ ppm): 2.32 (3H, s), 2.35 (3H, s), 4.13 (2H, s), 6.98 (1H, d, J = 8.2 Hz), 7.31 (1H, d, J = 8.4 Hz), 7.37 (3H, m), 7.42 - 7.47 (3H, m), 7.54 (1H, d, J = 8.4 Hz), 7.64 (2H, d, J = 7.8 Hz), 7.73 (1H, brs), 7.84 (2H, d, J = 8.1 Hz), 8.02 (1H, s), 11.36 (1H, s), 12.22 (1H, brs).

IR (Nujol): 1682 cm⁻¹.

m.p.: 150 - 154°C.

<Example 142; Production of 3-(4-bromo-2-chlorobenzyl)-2-methyl-5-(1-

pentanesulfonylcarbamoyl)indole (172)>

From 5-carboxy-3-(4-bromo-2-chlorobenzyl)-2-methylindole (0.15 g), N,N'-carbonyldiimidazole (0.128 g), diazabicycloundecene (0.12 g) and 1-pentanesulfonamide (0.12 g), obtained is 3-(4-bromo-2-chlorobenzyl)-2-methyl-5-(1-pentanesulfonylcarbamoyl)indole (172) (0.152 g).

$^1\text{H-NMR}$ (DMSO- d_6 , δ ppm): 0.82 (3H, t, $J = 7.2$ Hz), 1.24 - 1.32 (2H, m), 1.33 - 1.42 (2H, m), 1.67 - 1.72 (2H, m), 2.31 (3H, s), 3.48 (2H, t, $J = 7.7$ Hz), 4.07 (2H, s), 6.87 (1H, d, $J = 8.3$ Hz), 7.36 (1H, d, $J = 8.5$ Hz), 7.39 (1H, dd, $J = 1.9$ and 8.4 Hz), 7.64 (1H, dd, $J = 1.0$ and 8.4 Hz), 7.72 (1H, d, $J = 1.9$ Hz), 8.04 (1H, s), 11.39 (1H, s), 11.69 (1H, brs).

IR (Nujol): 1667 cm^{-1} .

m.p.: $168 - 172^\circ\text{C}$.

<Example 143; Production of 3-(4-bromo-2-chlorobenzyl)-2-methyl-5-(2-(5-chlorothieryl)sulfonylcarbamoyl)indole (173)>






From 5-carboxy-3-(4-bromo-2-chlorobenzyl)-2-methylindole (0.15 g), N,N'-carbonyldiimidazole (0.128 g), diazabicycloundecene (0.12 g) and 2-(5-chlorothieryl)sulfonamide (0.156 g), obtained is 3-(4-bromo-2-chlorobenzyl)-2-methyl-5-(2-(5-chlorothieryl)sulfonylcarbamoyl)indole (173) (0.146 g).

$^1\text{H-NMR}$ (DMSO- d_6 , δ ppm): 2.29 (3H, s), 4.07 (2H, s), 6.85 (1H, d, $J = 8.3$ Hz), 7.26 (1H, d, $J = 4.2$ Hz), 7.34 (1H, d, $J = 8.5$ Hz), 7.39 (1H, d, $J = 8.3$ Hz), 7.60 (1H, d, $J = 8.6$ Hz), 7.70 (1H, d, $J = 3.9$ Hz), 7.73 (1H, s), 8.00 (1H, s), 11.41 (1H, s), 12.51 (1H, brs).






IR (Nujol): 1689 cm^{-1} .

m.p.: $223 - 228^\circ\text{C}$.






1 / 29








2 / 29




3 / 2 9




4 / 29




5 / 29




(50)




(51)



(52)








(53)








(54)






6 / 2 9








7 / 2 9








8 / 29








9 / 29








10 / 29








11/29








12 / 29








13/29








14 / 29








15 / 29








16 / 29








17 / 29








18 / 29








19 / 29








20/29








21/29








22 / 29








23 / 29








24 / 29








25 / 29




26 / 29




27 / 29




28 / 29




(165)




(166)



(167)







(168)



(169)

29 / 29



It will be understood that various changes and modifications can be made in the details of procedure, formulation and use without departing from the spirit of the invention, especially as defined in the following claims.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
2217
2218
2219
2220
2221
2222
2223
2224
2225
2226
2227
2228
2229
2230
2231
2232
22